

Veterinary Package Insert

ERYTHROVET 30% W/W WATER SOLUBLE POWDER (30% Erythromycin Thiocyanate)

PRODUCT DESCRIPTION

White to off-white colour powder, which contains 300mg of erythromycin thiocyanate (277.6mg of erythromycin) per gram in 1kg sachet.

PHARMACODYNAMICS & PHARMACOKINETICS

Erythromycin is a macrolide antibiotic with a bacteriostatic action against a wide range of bacteria, although it also has a bactericidal action against certain pathogenic organisms.

Erythromycin exerts its effect only against multiplying bacteria. It inhibits protein synthesis by binding to 50 S ribosomal subunits of sensitive micro-organisms. Certain resistant micro-organisms with mutational changes in components of this subunit of the ribosome fail to bind the drug.

Erythromycin is absorbed rapidly when administered orally and diffusion occurs into most tissues and fluids. Elimination of erythromycin occurs primarily through hepatic metabolism and the remainder is excreted in active form in the urine and bile.

INDICATION

For the treatment of Chronic Respiratory Disease caused by Mycoplasma infection.

TARGETED SPECIES

Broilers up to 6 days of age.

Birds intended as replacement layers up to 6 days of age.

Broiler-Breeders of any age.

RECOMMENDED DOSAGE AND ADMINISTRATION

For oral administration via the drinking water.

Mix 59.45g of product per 64 litres of drinking water and administer continuously for 1-5 days, depending on response (approximately 25.5 mg/kg Erythromycin bodyweight per day).

Slowly add 59.45g to not less than 3.25 litres of clean, cold drinking water, stirring continuously until completely dissolved. This solution should then be added to more drinking water to make up a total volume of 64 litres.

CONTRAINDICATIONS

Any medicated drinking water which is not consumed within 24 hours should be discarded.

WARNINGS & PRECAUTIONS.

Special precautions for use in animals:

The water system to which the product is added must be in good working order and the header tanks and troughs must be free of dust, algae or other particulate matter.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Due to a lack of information on the fate and effects of erythromycin in the environment veterinary surgeons prescribing this product under the cascade should satisfy themselves that there will be no adverse impact on the environment.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not eat, drink or smoke while preparing the diluted formulation or while animals are being treated. In case of contact with eyes, rinse immediately with clean water and seek medical advice if irritation persists. Wash hands and exposed skin after use.

INTERACTIONS WITH OTHER MEDICATIONS

None reported.

PREGNANCY AND LACTATION

Not applicable.

ADVERSE EFFECTS

None reported.

OVERDOSE & TREATMENT

No information available.

WITHDRAWAL PERIOD

Chickens: 6 days.

Eggs: 6 days.

MAXIMUM RESIDUAL LIMIT (MRL)

Substance	Food	Maximum Residue Limits (MRLs) in µg/kg
Erythromycin	Edible offal, muscle, egg (mammalian and poultry)	300

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

500g

1kg

LIM SENG PHARMACHEM SDN. BHD.

(product registration holder)

2692, Tingkat Perusahaan Enam,

Kawasan Perusahaan Prai Mk 1,

13600 Prai, Pulau Pinang.

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.

Revised November 2019