



解熱鎮痛劑
解除疼痛及發燒
膜衣錠易吞服・無苦味

成分：
每錠含有 Acetaminophen 500 公絲。

作用：
普拿疼膜衣錠的止痛及解熱作用與水楊酸鹽相似。其止痛效果係來自周圍及中樞神經作用，而解熱作用則來自下視丘的調節中樞。

適應症：發燒、止痛（緩解頭痛、牙痛、咽喉痛、關節痛、神經痛、肌肉酸痛、月經痛）
普拿疼膜衣錠適用於解除各種疼痛，例如：頭痛、月經痛、肌肉骨節疼痛、肌肉痛及神經痛等。同時亦可用於伴有不適感及發燒之病症的止痛解熱，例如感冒及過熱性病。在牙科手術及拔牙後或在拔牙時，普拿疼膜衣錠均可作為有效的止痛劑。

副作用：

- 偶有過敏反應。在建議劑量下使用，少有副作用發生。
- 長期服用超大劑量會引起肝毒性，在兒童急性中毒劑量小於150mg/kg。

注意事項：

- 須置於小孩接觸不到之處。避免陽光直射，宜保存於陰涼之處。
 - 除非藥師、藥劑生或醫師指示，孕婦及授乳婦不建議自行使用。
 - 勿超過建議劑量，若有副作用產生，應立即停藥就醫。
- 警語：**
- 服用後，若有發疹、發紅、噁心、嘔吐、食慾不振、頭暈、耳鳴、喉嚨疼痛等症狀時，應停藥並就醫。
 - 除非藥師、藥劑生或醫師指示，服用本類藥品時，不得併服其他藥品或併服含酒精飲料。
 - 藥語：若你每天喝三杯或更多的酒精性飲料，你必須請教你的醫師是否能服用乙炔雌醇或其他類固醇劑，因乙炔雌醇可能造成肝損害。
 - 糖尿病（6-磷酸葡萄糖脫氫酶缺乏，Glucose-6-Phosphate Dehydrogenase Deficiency）患者不得服用本類藥品。
 - 除非有藥師、藥劑生或醫師指示，3歲以下不得服用此藥。
 - 曾經因藥物引起過過敏症者，使用前必須請教藥師、藥劑生或醫師。

用法用量：醫師或藥劑生指示藥品。

成人：每次 1-2 錠，每天 3-4 次。
孩童，7-12 歲：每次 1/2-1 錠，每天 3-4 次，最多可達 2 公克。

包裝：

6~1000 錠瓶裝、鋁箔盒裝。

施德隆股份有限公司製造 中華民國、台灣、台北
台北區淡水路146-7號

史克美占台灣分公司 總經理
地址：台北市信安路100號 電話：080-212259

64027906

Panadol® Film Coated Caplets
(Acetaminophen) 500mg

brand of **Acetaminophen**
For relief of pain and fever

Analgesic-antipyretic

Capsule-shaped Film Coated Caplets, easy to swallow.

Description

Each Film Coated Caplet contains Acetaminophen, 500mg.

Action

The analgesic and antipyretic actions of Acetaminophen are similar to those of the salicylates. Analgesia is mediated peripherally and also centrally, whereas antipyresis is produced by a central action on the hypothalamic regulatory center. Its micro-thin coating makes each 500mg caplet easy to swallow.

Indications

Panadol is indicated for the relief of painful disorders such as headache, dysmenorrhea, conditions involving musculoskeletal pain, myalgias and neuralgias. It is also indicated as an analgesic and antipyretic in conditions accompanied by discomfort and fever, such as the common cold and viral infections. Panadol is an effective analgesic after dental work and tooth extractions, and in teething.

Adverse Reaction

Allergic reactions occur occasionally. Following the recommended administration, adverse effects occur rarely. Prolonged severe overdose has been reported to have led to hepatotoxicity, children taking Acetaminophen 150mg/kg will induce acute toxicity.

Precaution

- Keep this product and all drug out of the reach of children.
- Store in sealed container at controlled temp. in dry place and keep from direct light.
- If you are pregnant or nursing a baby, seek the advice of a doctor or pharmacist before using this product.
- Do not take exceeding the recommended dosage, if adverse events occur, stop using this product immediately and go for a doctor.

Warning

- If skin rashes, nausea, vomiting, dysorexia, dizziness, tinnitus, sore throat occur, stop using this product immediately and go for a doctor.
- Do not take other drugs or alcoholic drinks when you are taking acetaminophen, unless directed by a doctor.
- If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers / fever reducers. Acetaminophen may cause liver damage.
- Favalism (Glucose-6-Phosphate Dehydrogenase Deficiency) do not take this product.
- Do not give this product to children under 3 years old, unless directed by a doctor or pharmacist.
- Seek the advice of a doctor or pharmacist before using this product, if you have ever been allergic to drugs.

Dosage and Administration

- Panadol is a doctor or pharmacist directed drug.
- Adults: 1-2 caplets, three or four times daily.
- Children, 7-12 years: 1/2-1 caplet, three or four times daily, up to 2 g.

How Supplied

Bottles or bottles of 6~1000 caplets.

Manufactured by

Sterling Drug (M) Sdn. Bhd.

Lot 89, Jalan Enggang, Ampang/Ulu Kelang Industrial Estate, 54200 Selangor

Usimulated by

SB SmithKline Beecham International INC., Taiwan R.O.C.

64027906