

TERNAFIN CREAM

DESCRIPTION:

A white to off-white, smooth homogenous mass.

COMPOSITION:

Each gram of cream contains: Terbinafine Hydrochloride (Micronised) 1% w/w.

ACTION AND PHARMACOLOGY:

Ternafin cream is a topical antifungal agent. It is an allylamine derivative with a broad spectrum of antifungal activity. It acts through inhibition of fungal sterol synthesis.

Pharmacodynamics:

Terbinafine is an allylamine which has a broad spectrum of antifungal activity. At low concentrations, terbinafine is fungicidal against dermatophytes, moulds and certain dimorphic fungi. Terbinafine is either fungicidal or fungistatic against yeasts, depending on the species. Terbinafine interferes specifically with fungal ergosterol biosynthesis at an early step. This leads to a deficiency in ergosterol and to intracellular accumulation of squalene, resulting in fungal cell death. Terbinafine acts by inhibiting squalene epoxidase in the fungal cell membrane. The enzyme squalene epoxidase is not linked to the cytochrome P-450 system. Terbinafine does not affect the metabolism of hormones or other drugs.

Pharmacokinetics:

Less than 5% of the dose is absorbed after topical application in humans. Systemic exposure is thus very slight. Biotransformation results in metabolites with no antifungal activities, which are excreted predominantly in the urine and the faeces. The half-life is 17 hours and there is no evidence of accumulation.

INDICATIONS:

Fungal infections of the skin caused by dermatophytes, eg. *Trichophyton* (eg, *T. rubrum*, *T. mentagrophytes*, *T. verrucosum*, *T. violaceum*), *Microsporum canis* and *Epidermophyton floccosum*; yeast infections of the skin, principally those caused by the genus *Candida* (eg, *Candida albicans*); *Pityriasis (tinea) versicolor* due to *Pityrosporum orbiculare* (also known as *Malassezia furfur*).

CONTRAINDICATIONS:

Hypersensitivity to terbinafine or any of the excipients contained in the cream.

SIDE EFFECTS:

Redness, itching or stinging occasionally occur at the site of application which does not necessarily require discontinuation. However, these reactions should be distinguished from allergic reactions which rarely occur but require discontinuation.

WARNING & PRECAUTIONS:

Ternafin cream is for external use only. Contact with the eyes should be avoided.

PREGNANCY & LACTATION:

Reproduction studies in animals do not suggest that there is any risk of foetal damage, but this has not been confirmed by controlled trials in pregnant women. Ternafin cream should therefore not to be administered during pregnancy unless the potential

benefits outweigh any potential risks. Terbinafine is excreted in breast milk and therefore its use in breastfeeding mothers cannot be recommended.

DRUG INTERACTIONS:

There are no known drug interactions

DOSAGE AND ADMINISTRATION:

To be applied once or twice daily. Cleanse and dry the affected areas thoroughly before application. Apply thin layer of cream to the affected skin and surrounding area and rub in lightly. In the case of intertriginous infections (submammary, interdigital, intergluteal, inguinal), the application may be covered with a gauze strip, especially at night.

Usual duration of treatment:

Tinea pedis, Tinea corporis, Tinea cruris: Apply twice daily for one week

Cutaneous Candidiasis: 1-2 weeks.

Pityriasis (tinea) versicolor: 2 weeks.

Relief of clinical symptoms usually occurs within a few days, but treatment must be used regularly and for an adequate length of time to avoid recurrence of infection. If there are no signs of improvement after 2 weeks, the diagnosis should be verified. Since experience with topical Ternafin Cream in children is limited, therefore it is not recommended to be used in children. While for the elderly, there is no evidence to suggest they require different dosages. Good general hygiene is necessary in conjunction with the use of Ternafin Cream in order to prevent re-infection (e.g. from underwear, socks, shoes, etc).

ROUTE OF ADMINISTRATION

Topical Administration

EFFECT ON ABILITY TO DRIVE AND USE MACHINE

None Known

OVERDOSAGE AND TREATMENT:

The adverse effects observed in man suggest that the main symptoms of an acute overdose would be gastrointestinal, eg. nausea or vomiting. If accidental ingestion of cream occurs, gastric lavage and/or symptomatic supportive treatment might then be required.

PRESENTATION:

Aluminium Tubes of 5, 10, 15, 50gm.

STORAGE CONDITIONS:

Store in a cool dry place below 30°C.

Protect from light and freezing.

Keep out of reach of children.

MANUFACTURED BY :

Noripharma Sdn. Bhd. 200701034604 (792633-A)
Lot 5030, Jalan Teratai
5 1/2 Mile off Jalan Meru
41050 Klang, Selangor, Malaysia

DATE OF REVISION:

April 2021