

DERMAFIN[®] CREAM

Ingredient(s):

Each gram contains:

Terbinafine Hydrochloride 10mg

Preservative(s):

Methyl Paraben 1mg

Propyl Paraben 1mg

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

Terbinafine is an allylamine with a range of antifungal activity its action being fungicidal against dermatophytes, moulds and certain dimorphic fungi, and either fungicidal or fungistatic against yeast, depending on the species.

Terbinafine interferes with fungal ergosterol biosynthesis at an early stage. This leads to a deficiency in ergosterol and to intracellular accumulation of squalene, resulting in fungal cell death. Terbinafine does not affect the metabolism of hormones or other drugs.

Indication(s):

Fungal infections of the skin caused by dermatophytes, eg. *Tricophyton*, *Microsporum canis* and *Epidermophyton floccosum*; yeast infections of the skin, principally those caused by the genus *Candida*; *Pityriasis (tinea) versicolor* due to *Pityrosporum orbiculare*.

Dosage and Administration:

Apply a thin layer and rub lightly to the affected skin and surrounding area once or twice daily. In the case of intertriginous infections (submammary, interdigital, intergluteal, inguinal), the application may be covered with gauze strip, especially at night.

Duration of treatment:

Tinea pedis, cruris, corporis: 1 week of treatment will normally suffice if the cream is applied once daily.

Cutaneous Candidiasis: 1 -2 weeks

Pityriasis (tinea) versicolor: 2 weeks

Contraindication(s):

Hypersensitivity to terbinafine.

Precaution(s) / Warning(s):

Terbinafine cream is for external use only. Contact with the eyes should be avoided. It is not recommended for pregnant and breastfeeding women

Drug Interaction(s):

There are no known drug interactions involving terbinafine cream.

Side Effect(s) / Adverse Reaction(s):

Occasional redness or itching.

Symptoms and Treatment for Overdosage, and Antidote(s):

No case of overdosage has been reported so far. The main symptoms of an acute overdosage may be gastrointestinal, eg. Nausea or vomiting. Gastric lavage and/or symptomatic supportive treatment might then be required.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container (Aluminium tube - air tight container).
Store at temperature below 30°C. Protect from light and moisture.

Product Description & Packing(s):

A white to off-white cream
Plastic jar of 500g (for export only).
Aluminium tube of 10g, 15g and 20g.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679