

B-MYCIN CREAM

Ingredient(s):

Each gram contains:

Betamethasone Dipropionate	0.643mg
(equivalent to Betamethasone 0.5mg)	
Gentamicin Sulphate (equivalent to Gentamicin	1mg)

Preservative(s):

Methyl Paraben	1mg
Propyl Paraben	1mg

Pharmacodynamic:

1. Betamethasone dipropionate topically produce prompt, marked and prolonged anti-inflammatory, antipruritic and vasoconstrictive effects. Gentamicin is bactericidal against a broad spectrum of common skin pathogens to which inflammatory skin disorders may be susceptible.
2. Excessive or prolonged use of topical corticosteroids can suppress pituitary- adrenal function resulting in secondary adrenal insufficiency and manifestations of hypercorticism. A single overdose of gentamicin would not be expected to produce symptoms. Excessive prolonged use of topical gentamicin may lead to overgrowth of lesions by fungi or nonsusceptible bacteria.

Pharmacokinetic:

1. Corticosteroids diffuse across cell membranes and complex with specific cytoplasmic receptors. These complexes then enter the cell nucleus, bind to DNA (chromatin), and stimulate transcription of messenger RNA (mRNA) and subsequent protein synthesis of various inhibitory enzymes responsible for the anti-inflammatory effects of topical corticosteroids.
2. Bacteria susceptible to the action of gentamicin include sensitive strains of *Streptococci*, *Staphylococcus aureus*, and the gram-negative bacteria *Pseudomonas aeruginosa*, *Aerobacter aerogenes*, *Escherichia coli*, *Proteus vulgaris* and *Klebsiella pneumoniae*.

Indication(s):

Treatment of impetigo, ecthyma and other localised primary bacterial skin infection with a gram-negative component and also for secondary bacterial infections complicating other pre-existing dermatoses.

Dosage and Administration:

Apply a thin layer to the affected area twice daily.

Mode of administration:

To be applied topically

Contraindication(s):

Hypersensitivity to any of B-mycin components.

Precaution(s) / Warning(s):

1. If irritation or sensitization develops with the use of B-mycin cream, treatment should be discontinued.
2. Since safety of topical corticosteroid use in pregnant women has not been established, drugs of this class should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.
3. B-mycin cream is not recommended for use in nursing mothers.
4. Pediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced hypothalamic-pituitary-adrenal (HPA) axis suppression and to exogenous corticosteroid effects than mature patients because of greater absorption due to a large skin surface area to body weight ratio.
5. HPA axis suppression, Cushing's syndrome, linear growth retardation, delayed weight gain and intracranial hypertension have been reported in children receiving topical corticosteroids. Manifestations

of adrenal suppression in children include low plasma cortisol levels and absence of response to ACTH stimulation. Manifestations of intracranial hypertension include a bulging fontanelle, headaches and bilateral papilledema.

6. Systemic absorption of topical corticosteroids will be increased if extensive body surface areas are treated or if the occlusive technique is used. Suitable precautions should be taken under these conditions or when long-term use is anticipated, particularly in infants and children.
7. Use of topical antibiotics occasionally allows overgrowth of nonsusceptible organism, including fungi. If this occurs or if irritation, sensitization or superinfection develops, treatment with gentamicin should be discontinued and appropriate therapy instituted.

Use of topical gentamicin preparations in closed hospital settings is actively discouraged.

Pregnancy & Lactation:

1. Since safety of topical corticosteroid use in pregnant women has not been established, drugs of this class should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.
2. B-mycin cream is not recommended for use in nursing mothers.

Side Effect(s) / Adverse Reaction(s):

Treatment with gentamicin has produced transient irritation (erythema and pruritus) that usually did not require discontinuance of treatment. Reported adverse reactions with the use of topical corticosteroids include: Burning, itching, irritation, dryness, hypopigmentation and allergic contact dermatitis.

Symptoms and Treatment for Overdosage, and Antidote(s):

Excessive or prolonged use of topical corticosteroids can suppress pituitary-adrenal function resulting in secondary adrenal insufficiency and manifestations of hypercorticism.

A single overdose of gentamicin would not be expected to produce symptoms.

Excessive prolonged use of topical gentamicin may lead to overgrowth of lesions by fungi or nonsusceptible bacteria.

Appropriate symptomatic treatment is indicated. Acute hypercorticism symptoms are virtually reversible. Treat electrolyte imbalance, if necessary. In cases of chronic toxicity, slow withdrawal of steroids is advised. Appropriate antifungal or antibacterial therapy is indicated if overgrowth occurs.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container (Aluminium tube- airtight container). Store at temperature below 30°C. Protect from light and moisture.

Product Description & Packing(s):

White cream.

Plastic jar of 500g (for export only).

Aluminium tube of 10g, 15g, 20g.



Manufacturer and Product Registration Holder :
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679