



KETOZOL SHAMPOO 2%

Description: Red coloured and bouquet fragrance shampoo.

Content:

Ketoconazole 2.0%w/w

Preservatives: methyl parabens 0.10%w/w; propyl parabens 0.01%w/w

Indication:

For the treatment and prophylaxis of pityriasis versicolor, seborrheic dermatitis and pityriasis capitis (dandruff).

Pharmacology:

Ketoconazole is an imidazole antifungal agent which interferes with ergosterol synthesis and therefore alters the permeability of the cell membrane of sensitive fungi. It is reported to be fungistatic at concentration achieved clinically. Ketoconazole has a wide spectrum of antimicrobial activity in vitro including activity against *Candida sp.*, *Blastomyces dermatitidis*, *Coccidioides immitis*, *Cryptococcus neoformans*, *Epidermophyton floccosum*, *Histoplasma capsulatum*, *Paracoccidioides brasiliensis*, *Malassezia furfur*, *Microsporum canis*, *Tricophyton mentagrophytes* and *T. rubrum*. As a shampoo, ketoconazole relieves scaling and pruritic conditions associated with pityriasis capitis (dandruff), seborrheic dermatitis and pityriasis versicolor. Systemic absorption following topical application in healthy subject is minimal.

Contraindications:

It is contraindicated in patients hypersensitive to ketoconazole.

Side effects/Adverse reactions:

After topical application, irritation, dermatitis or a burning sensation has occurred. In rare instances, mainly in patients with chemically damaged hair or grey hair, a discoloration of the hair has been observed.

Precautions:

Compliance with full course of therapy, fungal infection may require prolonged therapy. It should be applied for at least a few more days after the disappearance of symptoms. To prevent a rebound effect after stopping a prolonged treatment with topical corticosteroids, it is recommended to continue applying the topical corticosteroid together with Ketoconazole shampoo 2% and to subsequently and gradually withdraw the steroid therapy over a period of 2 - 3 weeks. Avoid shampoo to the eyes.

Pediatric use - safety and effectiveness in children have not been established.

Drug Interactions:

Nil for topical application.

Dosage and administration: (TOPICAL APPLICATION)

The affected areas of the skin or the hair should be washed with shampoo, which should be left on the skin for 3 - 5 minutes before rinsing.

Seborrheic dermatitis and pityriasis capitis (dandruff)

Treatment - twice weekly for 2 - 4 weeks. Prophylaxis - once every 1 - 2 weeks.

Pityriasis versicolor

Treatment - once daily for 5 days. Prophylaxis - once daily for 3 days for a single treatment course.

Treatment of overdosage:

Symptoms: Prolonged use of topical application of ketoconazole may rarely lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical use of the medication.

Treatment: Discontinue use and wash with water the affected part. For accidental ingestion, supportive measures should be carried out. To avoid aspiration, neither emesis or gastric lavage should be performed.

Storage conditions: Keep container tightly closed. Store in a cool, dry place below 30°C. Protect from light. Keep away from children.

Presentation: 60ml/bot. ; 250ml / bot.

Shelflife: 3 years.

Manufactured & : PRIME PHARMACEUTICAL SDN. BHD.

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.