

## **D3 Proposed Package Leaflet**

# **Biocillin 150mg/ml LA Suspension for Injection**

### **Active ingredient:**

Contains per ml:

Amoxicillin base (as amoxicillin trihydrate).....150mg

**Controlled Medicine**  
**Keep out of reach of children**  
**Jauhi dari kanak-kanak**  
**MAL registration number**

### **Product description:**

Off-white suspension.

### **Pharmacodynamics:**

Amoxicillin predominately inhibits cell wall synthesis in susceptible bacteria. Amoxicillin has a unique mode of action which directly and irreversibly disrupts existing cell wall peptidoglycan rather than newly forming peptidoglycan of the divisory septal wall as with other members of the penicillin family.

### **Pharmacokinetics:**

Amoxicillin is mainly distributed in the extracellular compartment. Its passage through the tissues is facilitated by its low degree of protein binding plasma levels (17%). The concentrations measured in lung tissue, pleural and bronchial are of the same order as plasma concentrations. Amoxicillin diffuses into the pleural and synovial fluid and into the lymphatic tissues. Amoxicillin is mainly excreted in urine - 10 to 25 % is excreted in the form of penicilloic acid. A part of it can also be excreted in bile. Metabolism is of little importance in elimination of amoxicillin. All penicillin's are eliminated rapidly after absorption.

### **Indication:**

Amoxicillin is effective against a wide range of Gram-positive and Gram-negative bacteria which include: *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus* species, *Salmonella* species, Staphylococci and Streptococci,

Not effective against beta-lactamase producing organisms.

Suitable for the control of infections due to susceptible microorganisms in cattle, sheep, pigs, dogs and cats where a single injection giving prolonged activity is required. It may also protect from secondary bacterial invasion due to sensitive organisms in cases where bacteria are not the initial cause of the disease. Indications include infections of:

- (a) Alimentary tract
- (b) Respiratory tract
- (c) Skin and soft tissue
- (d) Urogenital tract and,
- (e) In prevention of post-operative infection (treat before surgery).

### **Recommended dose:**

To ensure the correct dosage, body weight should be determined as accurately as possible to avoid underdosing.

Cattle, sheep and pigs: by intramuscular administration.

Dogs and cats: - by subcutaneous injection.

General: 15 mg per kg body weight, equivalent to 1.0 ml per 10 kg body weight.

This dosage can be repeated after 48 hours when necessary. Shake the vial before use

Swab the septum before removing each dose. Use a dry sterile needle and syringe. A separate injection site should be used for each administration.

| <b>Animal</b> | <b>Weight (kg)</b> | <b>Dosage volume (ml)</b> |
|---------------|--------------------|---------------------------|
| Cattle        | 450 kg             | 45.0 ml                   |
| Sheep         | 65 kg              | 6.5 ml                    |
| Pigs          | 150 kg             | 15.0 ml                   |
| Dogs          | 20 kg              | 2.0 ml                    |
| Cats          | 5 kg               | 0.5 ml                    |

Dose volume is equivalent to 1 ml per 10 kg bodyweight. If dose volume exceeds 20 ml, it should be divided and injected into two sites. As with other injectable preparations normal aseptic precautions should be observed.

An appropriately graduated syringe must be used to allow accurate administration of the required dose volume. This is particularly important when injecting small volumes.

**Contraindication:**

This product is not suitable for intravenous or intrathecal use. This product should not be administered to rabbits, hamsters, gerbils or guinea pigs. Not for use in known cases of hypersensitivity to penicillins or cephalosporins

**Warning and Precautions:**

**Special warnings for each target species**

No special warnings.

**Special precautions for use**

**Special precautions for use in animals**

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local epidemiological information.

**Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Care should be taken to avoid accidental self-injection. In the case of accidental self-injection, seek medical advice immediately.

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.

Handle this product with great care to avoid exposure, taking all recommended precautions.

If you develop symptoms following exposure such as skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing, are more serious symptoms and require urgent medical attention.

Wash hands after use.

**Interaction of other medicaments:**

It is not generally recommended to use bactericidal and bacteriostatic antibiotics at the same time.

**Pregnancy and Lactation:**

Can be safely administered during pregnancy and lactation.

**Side Effects:**

Use of the product may occasionally result in local tissue reaction.

**Symptoms and treatment of overdose:**

Penicillins have a wide safety margin.

**Withdrawal period:**

Cattle:

Meat and offal: 28 days

Milk: 84 hours

Pigs:

Meat and offal: 19 days

Sheep:

Meat and offal: 19 days

Not for use in sheep producing milk for human consumption.

**Storage condition:**

Store below 30°C. Protect from light.

**Packaging:**

Cardboard box with a colourless type II glass bottle of 50 and 100 ml, closed with a sterile rubber stopper and aluminium cap.

**Manufacturer:**

Interchemie werken "De Adelaar" Eesti AS  
Vanapere tee 14, Püüsi village, Viimsi municipality, Harju country, 74013 Estonia.

**Marketing Authorization Holder:**

Danberg (M) Sdn Bhd  
Lot 1, Jalan Bursa 23/4, Seksyen 23, 40300 Shah Alam, Selangor.  
Tel: 603-5548 4905

Date of revision: 28 January 2020