

MOXIFEED 500G/KG GRANULES

Description and Composition

MOXIFEED 500G/KG GRANULES is an off white to yellowish granules. Each kg contains 500g Amoxicillin Trihydrate as its active ingredient.

Pharmacodynamics

Amoxicillin is semi-synthetic penicillin, derived from the 6 APA nucleus (6-penicillanic acid). It is a slightly acidic, weakly liposoluble, acid-stable compound which can be administered enterally or parenterally. Amoxicillin is a broad-spectrum antibiotic. It causes alterations of the bacterial wall inducing secondarily an osmotic lysis of the bacterium. Amoxicillin has a predominantly bactericidal effect on Gram-positive and Gram-negative bacteria, including *Staphylococcus* spp., *Streptococcus* spp., *Corynebacterium* spp., *Escherichia* spp., *Haemophilus* spp., *Klebsiella* spp., *Shigella* spp., *Salmonella* spp., *Proteus* spp., and *Pasteurella* spp.

Pharmacokinetics

Amoxicillin is well absorbed after oral administration. The fraction reaching the bloodstream is poorly metabolized, the metabolite identified in the laboratory species being penicillic acid, which is rapidly eliminated in the urine. Amoxicillin is weakly bound to plasma proteins (about 20%) and is widely distributed to different organs.

Indication

Broiler chickens: For the prevention of respiratory infections caused by *E. coli* susceptible to Amoxicillin.

Recommended Dose

Broiler chickens: 20mg Amoxicillin equivalent to 23mg Amoxicillin trihydrate (46mg Moxifeed 500g/kg Granules) per kg body weight per day for 5 consecutive days in feed.

Route of Administration

For oral administration via feed.

Contraindication

Do not use in animal known for their sensitivity to penicillins or other beta-lactams. Do not use in guinea pigs, rabbits, hamsters and gerbils.

Warning and Precautions

Special warnings for each target species

None known.

Special precautions for use

Special precautions for use in animals

This medicated premix is intended for the manufacture of medicated feed and cannot be used as it is; the rate of incorporation of the medicated premix into the food may not be less than 5kg/tonne.

Special precautions to be taken by the person administering the veterinary medicinal product to animals
Penicillins and cephalosporins may cause a hypersensitivity reaction following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may cause cross-reactivity with cephalosporins and vice versa. The allergic reactions to these substances can sometimes be serious. Do not handle this product if you have a history of allergy to cephalosporins and penicillins, or if you have been advised against working with such preparations. Handle this product with the recommended precautions to avoid any risk of exposure: Wearing masks, goggles and protective gloves is recommended when preparing the feed.

If you experience symptoms such as rash after exposure, consult your doctor and show this warning. Swelling in the face, lips or eyelids, or breathing difficulties are more serious symptoms and require urgent medical attention.

Other precautions

None.

Interaction with Other Medicaments

Not known.

Pregnancy and Lactation

See section "Withdrawal Periods".

Side Effects

Not known.

Symptoms and Treatment of Overdose

No adverse effects were observed after administration of the drug at doses 5 times higher than recommended.

Storage Condition

Store below 30°C.

Shelf Life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Periods

Meat and offal:

Chickens: 6 days

Eggs: In the absence of MRLs for eggs, do not use in laying hens producing eggs for human consumption (4 weeks before start and during egg production).

Packing

10kg, 25kg.

Revised date: 6.3.2026

	Manufacturer & Product Registration Holder:
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com