

Amogen

Suspension for injection

COMPOSITION: Each ml contains:

Amoxicillin trihydrate	150 mg
Gentamicin Sulfate	40,000 IU
Benzyl alcohol (as preservative)	0.03 ml

DESCRIPTION

The finished product is ivory cream suspension.

PHARMACODYNAMICS

Amoxicillin is a bactericidal antibiotic that belongs to the penicillin group. It inhibits the synthesis of the mucopeptide, one of the components of the bacterial rigid cell wall that gives the bacteria mechanical stability, destroying the bacteria in its growth phase.

Gentamicin is an aminoglycoside antibiotic that acts directly on the ribosome, where it inhibits protein biosynthesis and decreases the fidelity of interpretation of the genetic code. The penetration of gentamicin in the cell is increased when the permeability of the bacterial cell wall is altered by amoxicillin. This fact allows the use of smaller amounts of the aminoglycoside.

The bound amoxicillin and gentamicin have a broad antibacterial spectrum that includes Gram positive bacteria (*corinebacteria*) and Gram negative bacteria (*E. coli*, *Pasteurella spp.*, and *Salmonella spp.*). Their association improves the therapeutic properties of each of these antibiotics administered alone.

PHARMACOKINETICS

Amoxicillin is well absorbed after parenteral administration. 17-20% of the active substance binds to plasma proteins. It is excreted mainly by unaltered urine and to a lesser extent by milk and bile.

Gentamicin is poorly absorbed from the

gastrointestinal tract, however, it is well absorbed after parenteral administration. It is rapidly distributed reaching the kidney, liver, lung, endometrium and breast parenchyma. Its elimination takes place in a biphasic manner, a first rapid phase corresponding to the distribution of the active substance in the tissues and a second slow phase corresponding to the release of the active substance from the tissues to the plasma and its subsequent elimination from the plasma.

INDICATION

Cattle: Pneumonia caused by *Pasteurella multocida* and *Arcanobacterium pyogenes* strains sensitive **.

Diarrhea caused by strains of *Escherichia coli* and *Salmonella spp* sensitive **.

Pigs: Pneumonia caused by strains of *Pasteurella multocida*, *Haemophilus parasuis* and *Actinobacillus pleuropneumoniae* sensitive **.

Colibacillosis Produced by sensitive strains **.

** Sensitive to amoxicillin and gentamicin.

TARGET SPECIES

Cattle and pigs.

ROUTE OF ADMINISTRATION

Intramuscular route

RECOMMENDED DOSE

Cattle and pigs: 15 mg amoxicillin trihydrate and 4,000 IU gentamicin sulfate/ kg bw equivalent to 1 ml/ 10 kg body weight for 3 days.

CONTRA-INDICATION

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

Do not use in animals hypersensitive to penicillin. Do not use in animals with impaired renal function. Do not use in pregnant animals.

INTERACTION WITH OTHER MEDICAL PRODUCT

It should not be used together with bacteriostatic antibiotics.

WARNINGS AND PRECAUTIONS

Special warnings for each target species

None

Special precautions for use in animals

Shake the product vigorously before using. When the vial has been at rest for a while, separation into two layers that are easily resuspended by agitation is normal.

In the administration of high dosages, it is advised to divide the medicine between two or more injection points, in order to favor absorption.

Special precautions to be taken by the person administering the medicinal product to animals

In case of accidental self-injection, see a doctor and show the package leaflet or label.

People with known hypersensitivity to gentamicin and / or penicillin should avoid all contact with the veterinary medicinal product.

-Keep out of reach of children

-Jauhkan daripada kanak-kanak

PREGNANCY AND LACTATION

The product crosses the placental barrier and can cause ototoxicity and nephrotoxicity in the fetus. Its use is not recommended during pregnancy.

SIDE EFFECTS

Very rarely, in prolonged treatment with doses higher than those recommended, renal dysfunction may occur.

Very rarely, hypersensitivity reactions occur in

sensitive animals.

The frequency of adverse reactions should be classified according to the following groups:

- Very frequently (more than 1 animal for every 10 animals treated has adverse reactions)
- Frequently (more than 1 but less than 10 animals per 100 treated animals)
- Infrequently (more than 1 but less than 10 animals per 1,000 treated animals)
- Rarely (more than 1 but less than 10 animals per 10,000 treated animals)
- Very rarely (less than 1 animal per 10,000 treated animals, including isolated cases).

SYMPTOMS AND TREATMENT OF OVERDOSE

In case of poisoning due to overdosing, discontinue the medication and administer symptomatic treatment. The administration of the medicine in pigs and cattle, at triple doses of the recommended one and for a time three times higher than the recommended cause evident lesions in the administration area; In pigs, alterations related to hepatitis or nephritis may occur.

STORAGE CONDITION

Store in a dry place not exceeding 30°C.

SHELF LIFE: 2 years

SHELF LIFE AFTER OPENING: 24 hours

WITHDRAWAL PERIOD

Cattle (Meat): 140 days.

Pigs (Meat): 84 days.

Not permitted for use in animals whose milk is used for human consumption.

Do not use in pregnant females whose milk is used for human consumption in the 2 months prior to the expected date for delivery.

PACKAGING: 100 ml

REVISED DATE: Mar 17, 2020