

Amox Excel 500mg/g Water Soluble Powder

For Veterinary Use Only

Composition:

Each gram contains:

Amoxicillin trihydrate: 500mg

Product description:

A white or almost white powder.

Pharmacodynamics:

Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against Gram positive and Gram negative bacteria. IT owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

Pharmacokinetics:

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

Indication:

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin.

Not effective against beta-lactamase producing organisms.

Use of the product should be based on susceptibility testing and it should take into account official and local antimicrobial policies.

Contraindications:

Should not be administered to rabbits, hamsters, gerbils and guinea pigs, or to birds producing eggs intended for human consumption.

Interaction with other medicaments:

None known.

Symptoms and treatment of overdose:

No problems with overdosage have been reported. Treatment should be symptomatic and no specific antidote is available.

Side effects:

None known.

Recommended dose:

Chickens, turkeys and ducks:

The product is administered in the drinking water. Prepare the solution with fresh tap water immediately before use. Any unused medicated water should be discarded after 24 hours.

In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated.

The following formula may be used to calculate the amount of product required per day (in grams):

$$\frac{\text{Number of birds} \times \text{average live weight (kg)}}{25 \text{ (for 20 mg/kg) or } 33 \text{ (for 15 mg/kg)}}$$

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing. The uptake of medicated water depends on the clinical condition of the birds. In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted taking into account water intake.

Chickens:

The recommended dosage is 15 mg amoxicillin trihydrate (30 mg Amox Excel 500mg/g Water Soluble Powder) per kg body weight. The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks:

Recommended dosage is 20 mg amoxicillin trihydrate (40 mg Amox Excel 500mg/g Water Soluble Powder) per kg bodyweight for 3 consecutive days.

Turkeys:

Recommended dosage is 15-20 mg amoxicillin trihydrate (30-40 mg Amox Excel 500mg/g Water Soluble Powder) /kg bodyweight for 3 days or in severe cases for 5 days.

Pregnancy and lactation:

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.

Use only according to the benefit/risk assessment of the responsible veterinarian.

Withdrawal period:

Chickens (meat & offal): 1 day

Ducks (meat & offal): 9 days

Turkeys (meat & offal): 5 days

Not for use in laying birds producing eggs for human consumption

Packaging:

100g aluminium pouch

Warning and precautions:

Special warnings for each target species

None.

Special precautions for use in animals

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact.

Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1) Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.

2) Handle this product with great care to avoid exposure, taking all recommended precautions.

3) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

- Avoid inhalation of dust. Wear either a disposable half-mask respirator or a non disposable respirator
- Wear gloves during preparation and administration of medicated water or liquid feed
- Wash any exposed skin after handling the product or medicated water or feed
- Wash hands after use

Storage condition:

Store below 30°C.

Shelf life: 36 months

Shelf life after first opening of container: 7 days

Shelf life after reconstitution/dilution: 24 hours

Manufacturer:

Nam Pharma SdnBhd
5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak.

Marketing Authorization Holder:

F.E Venture SdnBhd
Lot 3 & 5, Jalan PJS 11/8,
Bandar Sunway, 46150 Petaling Jaya,
Selangor, Malaysia