

PACKAGE INSERT (FRONT)

hevid

TRISILON CREAM

VIVIR02-2

DESCRIPTION

White, opaque cream

COMPOSITION

Betamethasone Dipropionate equivalent to
Betamethasone 0,05% w/w
Clotrimazole 1,00% w/w
Gentamicin Sulfate equivalent to Gentamicin 0,10% w/w

ACTIONS AND PHARMACOLOGY

Trisilon Cream is a combined preparation of the anti-inflammatory, antipruritic and vasoconstrictive actions of betamethasone dipropionate with the wide-spectrum antifungal activity of clotrimazole and the broad-spectrum bactericidal antibiotic activity of gentamicin sulfate.

Betamethasone dipropionate, a corticosteroid, diffuses across cell membranes, binds with DNA, and stimulates transcription of messenger RNA (mRNA) resulting in synthesis of various enzymes responsible for the anti-inflammatory effects of betamethasone.

Clotrimazole, an imidazole antifungal, appears to act on the fungal cell membrane, causing leakage of cell contents.

Gentamicin, an aminoglycoside and a bactericidal, provides highly effective topical treatment in primary and secondary bacterial infection of the skin. It is effective against many gram-negative aerobes and against some strains of staphylococci.

PHARMACOKINETICS

Betamethasone can be absorbed when given locally to give systemic effects. Clotrimazole can penetrate intact skin but there is little if any systemic absorption. Gentamicin has been reported to give systemic absorption after topical use. Gentamicin do not appear to be metabolized and are excreted virtually unchanged in the urine.

INDICATIONS

For the relief of the inflammatory manifestations of corticosteroid-responsive dermatoses when complicated with secondary infection caused by bacteria (sensitive to gentamicin) and fungi (sensitive to clotrimazole) or when the possibility of such infections is suspected.

Effective in treating tinea pedis, tinea cruris and tinea corporis caused by *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Epidermophyton floccosum* and *Microsporum canis*, candidiasis due to *Candida albicans* and tinea versicolor caused by *Malassezia furfur* (*Pityrosporum obiculare*).

Used in the treatment of severe systemic infections due to sensitive strains of *Staphylococcus aureus* (coagulase positive, coagulase negative and some penicillinase-producing strains) and the gram-negative bacteria: *Pseudomonas aeruginosa*, *Aerobacter aerogenes*, *Escherichia coli*, *Proteus vulgaris* and *Klebsiella pneumonia*.

CONTRAINDICATIONS

Trisilon Cream is contraindicated in patients with hypersensitivity to any of the ingredients.

PREGNANCY AND LACTATION

Safety for use in pregnancy and lactation has not been established. Drugs of this class should be used during pregnancy only if the potential benefit outweighs the potential risk to the fetus. Administration should not be used extensively in large amounts or for prolonged periods in pregnant women.

PRECAUTIONS

- Cross-allergenicity among aminoglycosides has been observed.
- Treatment should be discontinued if irritation or sensitization develops.
- Avoid use on face, groin, or in the axillae.

XX

PACKAGE INSERT (BACK)

hovid

- Side effects that are reported following systemic use of corticosteroids may occur with topical corticosteroid, especially in infants and children.
- Use of topical gentamicin preparations in closed hospital settings is actively discouraged. Application to open wounds or damaged skin should be avoided.
- Long term treatment of gentamicin is not advised. Prolonged usage of topical antibiotics may result in overgrowth of nonsusceptible microorganisms. Treatment should be discontinued if this situation arises.
- Caution should be exercised when the product is used to treat extensive body surface areas, or if occlusive dressings are used especially for prolonged periods, or in the presence of dermal disruption.

INTERACTION WITH OTHER MEDICAMENTS AND OTHER FORMS OF INTERACTION

Not applicable.

MAIN SIDE/ADVERSE EFFECTS

Adverse reactions have been very rarely reported including skin discolouration, hypochromia, burning, erythema, exudation and pruritus.

The following local adverse reactions have been observed with the topical usage of:

Corticosteroids (especially under occlusive dressing): burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruption, hypopigmentation, perioral dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, skin atrophy, striae and miliaria.

Clotrimazole: erythema, stinging, blistering, peeling, edema, pruritus, urticaria and general irritation of the skin.

Gentamicin: transient irritation (erythema and pruritus) that usually did not require discontinuance of treatment.

TRISILON CREAM

OVERDOSE

Symptoms

Excessive or prolonged application of topical corticosteroids can suppress pituitary-adrenal function, resulting in secondary adrenal insufficiency, and produced hypercorticism, including Cushing's disease.

Overdosage by topical clotrimazole application is highly improbable.

Single overdosage of gentamicin would not be expected to produce symptoms. Excessive or prolonged use of topical antibiotics may lead to overgrowth of lesions by nonsusceptible microorganisms.

Treatments

In the event of chronic toxicity, slow withdrawal of corticosteroid is recommended. Acute hypercorticotid symptoms are generally reversible.

DOSAGE AND ADMINISTRATION

Apply a thin film over the affected and surrounding skin areas

Adults: 2 times daily

Apply regularly for effective treatment. Duration of therapy varies according to the extent and location of disease and patient response.

Storage: Store below 30°C

Presentation / Packing: Cream 15 g

Manufactured by: HOVID Bhd,
121, Jalan Tunku Abdul Rahman, 30010 Ipoh, Malaysia.

Date of information: July 2013.

hovid