

Veterinary Package Insert

KHC Mebendazole 60MG/G Premix Powder

PRODUCT DESCRIPTION

White or almost white colour powder, which contains 60mg of mebendazole per gram in 500gm or 1kg sachet.

PHARMACODYNAMICS & PHARMACOKINETICS

Mebendazole is an anthelmintic belonging to the benzimidazole group of compounds with efficacy against certain gastrointestinal roundworms and lungworms in horses.

Mebendazole acts selectively against the gastrointestinal and lung parasites in the host. It is based on an interaction with the microtubular system of the absorptive cells of the worm, leading to an irreversible lytic necrosis of those cells and death of the worm.

The pharmacokinetic profile of mebendazole is similar in various animal species, including horses. Mebendazole has a poor oral bioavailability, due to a low solubility in aqueous systems, a slow dissolution rate in the gastrointestinal tract and first-pass metabolism in the gut wall and the liver. This causes a high faecal excretion of parent drug and low levels in plasma and tissues. Absorption is not linearly dependent of dose. Highest concentrations of mebendazole-related residues are found in the liver and kidneys and these consist mainly of metabolites. The biotransformation of mebendazole involves carbamate hydrolysis, ketone reduction and conjugation. The elimination from plasma and tissues is rapid, although there is some retention of residues in the liver.

The systemic bioavailability of mebendazole in horses is very low, irrespective of the oral dosage form. After a 6.5 mg/kg dose, the concentrations in plasma never exceed 10 ng/ml.

INDICATION

An oral broad spectrum anthelmintic for the treatment of helminthiasis in the horse.

The product is effective against benzimidazole - susceptible strains of the following worms:

Large strongyles:	<i>Strongylus vulgaris</i> <i>Strongylus edentatus</i> <i>Strongylus equinus</i> <i>Triodontophorus</i> spp.
Small strongyles:	<i>Cyathostomes</i>
Stomach worms:	<i>Trichostrongylus axei</i>

Ascarids:	<i>Parascaris equorum</i>
Pinworms:	<i>Oxyuris equi</i> <i>Probstmayria vivipara</i>
Lungworms:	<i>Dictyocaulus arnfieldi</i>

RECOMMENDED DOSAGE AND ADMINISTRATION

Add to feed immediately before administration.

Discard any remaining medicated feed.

To ensure administration of a correct dose, bodyweight should be determined as accurately as possible.

Horses:

The dose rate is 5-10 mg mebendazole per kg bodyweight orally.

Repeat this dose every 6 weeks.

Horses 200-400 kg 33.33g of product

Horses 401-800 kg 66.67 g of product

CONTRAINDICATIONS

None.

WARNINGS & PRECAUTIONS.

Special warnings for each target species:

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy.

- Too frequent and repeated use of anthelmintics from the same class, over an extended period of time.
- Underdosing, which may be due to underestimation of bodyweight, misadministration of the product, or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

The use of this product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics.

To reduce this risk dosing programmes should be discussed with your veterinary surgeon.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wash hands after use

Persons with known hypersensitivity to ingredients in the product should not handle the product.

Avoid contact with skin and eyes. Wash off any contact with the product immediately with clean running water. Seek medical attention if irritation persists.

INTERACTIONS WITH OTHER MEDICATIONS

No data available.

PREGNANCY AND LACTATION

Animals should not be treated with a dose of 15 mg/kg or over during the first 4 months of pregnancy.

ADVERSE EFFECTS

None reported.

OVERDOSE & TREATMENT

Supportive therapy if required.

WITHDRAWAL PERIOD

Not to be used in horses intended for human consumption.

Treated horses may never be slaughtered for human consumption.

MAXIMUM RESIDUAL LIMIT (MRL)

Pharmacologically active substance(s)	Target Tissues	Maximum Residue Limits (MRLs) in µg/kg
Mebendazole	Muscle, liver (Equidae)	60
	Fat (Equidae)	60
	Liver (Equidae)	400
	Kidney (Equidae)	60

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

500g

1kg

KHENG HONG CHAN FARMING SDN. BHD.

(product registration holder)
No.49 Lorong Alma Jaya 12,
Kawasan Industri Alma,
14000 Bukit Mertajam,
Pulau Pinang.

THYE PHARMA SDN. BHD.

(manufacturer)
No.2 Lorong Industri Ringan Permatang
Tinggi 8,
Kawasan Industri Ringan Permatang Tinggi,
14000 Bukit Mertajam,
Pulau Pinang.

Revised August 2016