

Oorafax 15%w/w
Powder
(FOR VETERINARY USE ONLY)

COMPOSITION

Chlortetracycline Hydrochloride 150 mg/g

PRODUCT DESCRIPTION

The finished product is a yellow color powder.

TARGET SPECIES

Pigs, chickens, turkeys or ducks

PHARMACODYNAMIC

Chlortetracycline is a natural broad spectrum antibiotic of the tetracycline group. Its activity is bacteriostatic by interfering with the protein synthesis at the S30 ribosome sub-unit. At recommended dosages it has no pharmacological effects on cardiovascular, nervous or other body systems.

PHARMACOKINETIC

When dosed orally it is quickly absorbed into the blood stream. In-feed dosing provides steady to near steady state plasma levels. This provides effective concentrations in various tissues, including lungs and other respiratory tissues. The major metabolite is 4-epi-chlortetracycline.

INDICATIONS

For treatment and control of respiratory and systemic infections associated with organisms sensitive to chlortetracycline.

WARNINGS AND PRECAUTIONS

Special warnings for each target species:

None

Special precautions for use:

i. Special precautions for use in animals:

Official, national and regional antimicrobial policies should be taken into account when the product is used.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals:

If you know you are hypersensitive (allergic) to chlortetracycline, do not handle the product.

When handling this product, inhalation of the dust must be avoided by wearing a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143. Avoid contact with skin and eyes. Gloves should be worn whilst handling this product. If

contact with skin or eyes occurs, wash area immediately with clean fresh water. If irritation persists seek medical attention. Do not eat, drink or smoke whilst handling the product. Hands and exposed skin should be washed thoroughly after use.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

CONTRAINDICATIONS

Do not use in ruminants.

ADMINISTRATION & DOSAGE

Premix for medicated feeding stuff

The recommended dose rates are as follows:

Pigs: 10–20 mg/kg bodyweight daily
Chickens: 20–30 mg/kg bodyweight daily
Turkeys: 10–30 mg/kg bodyweight daily
Ducks: 10–30 mg/kg bodyweight daily

These dose rates can usually be achieved by mixing 2.0 kg of Orafax 15%w/w Powder per ton of complete feed to give a concentration of 300 ppm chlortetracycline hydrochloride. However, the correct incorporation rate should always be calculated based on the feed consumption rates of the animals to be treated.

The intake of medicated feed depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of chlortetracycline has to be adjusted accordingly.

To ensure thorough dispersion of the product, it should first be mixed with a suitable quantity of feed ingredients before incorporation into the final mix.

The medicated feed should be supplied to the affected pen(s) or group(s) of pigs, chickens, turkeys or ducks. Treatment should be continued for a period of five to seven days for respiratory or systemic infections. During the treatment period, only feed medicated with the product should be supplied.

For incorporation into dry feed at the registered mill. A manufacturer who is approved to incorporate directly at any concentration, veterinary medicinal products or pre-mixtures containing such products must be responsible for mixing when incorporation is less than 1.33 kg per ton for final feed.

INTERACTION WITH OTHER MEDICAMENTS

Chlortetracycline is sometimes used in association with other antimicrobials, notably sulphadimidine, penicillin and neomycin. There are no known adverse reactions with these or any other substances.

PREGNANCY AND LACTATION

No problems known

OVERDOSE AND TREATMENT

Chlortetracycline is of low toxicity and there is a wide safety margin at the recommended dosage. On rare occasions overdosage may cause diarrhea and over growth of yeast and fungi. Under such conditions, withdraw medication and apply appropriate treatment.

SIDE EFFECT

The product is of low toxicity and side effects are rarely encountered. If suspected adverse reactions occur, treatment should be discontinued immediately.

WITHDRAWAL TIME

Meat: Animals must not be slaughtered for human consumption during treatment.

Pigs: 10 days

Chickens and Turkeys: 2 days

Ducks: 4 days

Eggs: (Chickens, Turkeys and Ducks) Eggs of treated animals should not be used for human consumption.

SHELF LIFE: 36 months

SHELF LIFE after first opening & reconstitution: 24 hours

STORAGE

Store at temperature not exceeding 30°C. Strictly avoid light & heat exposure.

PACKAGING

The powder filled 25 kg in multi-wall bag.

DATE OF REVISION: 09 May 2019



Product Registration Holder & Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN BHD
神農動物保健(馬)有限公司
No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,
Taman Perindustrian USJ 1,
47610 Subang Jaya, Selangor, Malaysia.



Distributed By:
ZAGRO CHEMICALS SDN. BHD.
No. 27, Jalan PJS 3/34,
Taman Sri Manja,
46000 Petaling Jaya, Selangor, Malaysia.