

TETRA 20%w/w

POWDER
(FOR VETERINARY USE ONLY)

COMPOSITION

Oxytetracycline 200 mg/g

PRODUCT DESCRIPTION

The finished product is a light yellowish or brownish color powder.

TARGET SPECIES

Sheep (lambs), goats (young goats), pigs, poultry (chickens, turkeys, ducks)

PHARMACODYNAMIC

Oxytetracycline binds reversibly to receptors of the 30S ribosomal fraction, this leads to a blocking of the binding of aminoacyl-tRNA to the corresponding site of the ribosome complex messenger-RNA. This results in inhibition of protein synthesis and therefore a growth arrest of the bacterial culture. Oxytetracycline is a bacteriostatic. The bacteriostatic activity of oxytetracycline involves penetration of the substance into the bacterial cell. The penetration of oxytetracycline is exercised by both passive and active dissemination. The main possible resistance mode is related to the presence of an R factor responsible for a decrease in active transportation oxytetracycline. Oxytetracycline is a broad spectrum antibiotic. It is primarily active against microorganisms Gram positive and negative, aerobic and anaerobic, and against mycoplasma, Chlamydia and Rickettsiae. The acquired resistance to oxytetracycline was reported. Such resistance is usually plasmid origin. Crossresistance to other tetracyclines is possible. Continuous treatment with low doses of oxytetracycline can also lead to increased resistance to other antibiotics.

PHARMACOKINETIC

For the majority of species, oxytetracycline is quickly (24 hours) absorbed after oral administration in animals fasted and its bioavailability is between 60% and 80%. The bioavailability may decrease in the presence of food in the stomach because oxytetracycline forms insoluble chelates with divalent or trivalent cations (Mg, Fe, Al, Ca) they contain. In pigs, the influence the food is negligible bioavailability of oxytetracycline which is less than 5%. Oxytetracycline binds to plasma proteins variably depending on the species (20-40%). Its distribution is wide. Oxytetracycline diffuses throughout the body, the highest concentrations found in the kidneys, liver, spleen and lungs. Oxytetracycline crosses the placenta. Oxytetracycline is excreted unchanged mainly via urine. It is also excreted in the bile, but a large percentage of oxytetracycline is reabsorbed by the small intestine (enterohepatic cycle).

INDICATIONS

In lambs, young goats, pigs, chickens, turkeys and ducks: Treatment and infected based prevention of respiratory infections and infections digestive caused by germs sensitive to oxytetracycline.

WARNINGS AND PRECAUTIONS

Special warnings for each target species:

None

Special precautions for use:

i. Special precautions for use in animals:

This premix is intended for the manufacture of medicated feed and cannot be used as such; The incorporation rate of the premix to the feed can be less than 5 kg / ton.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not handle this product in case of known allergy to tetracyclines.

If reaction after exposure to the product (e.g. rash), consult a doctor.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

CONTRAINDICATIONS

Do not use in cases of known allergy to oxytetracycline or other substance tetracycline.

Do not use in case of known resistance to tetracyclines.

ADMINISTRATION & DOSAGE

Oral route

Pigs: Prevention: 25 mg oxytetracycline per kg body weight per day, orally, for 10 days. Treatment: 50 mg oxytetracycline per kg body weight per day, orally, for 10 days.

Lambs, young goats and poultry (chickens, turkeys & ducks): Prevention: 20 mg oxytetracycline per kg bodyweight per day, orally, for 10 days. Treatment: 40 mg oxytetracycline per kg body weight per day, orally, for 10 days.

The amount of food consumed drug by animals depends on their physiological state (age) and clinical. To meet the recommended doses oxytetracycline and consider the food ingested workforce, the incorporation rate can be increased or possibly reduced to the minimum incorporation limit of 5 kg / ton. This can lead to different ppm oxytetracycline levels in medicated feed those mentioned below indication.

Poultry, lambs and young goats: For daily feed intake of 20 g of feed per kg bodyweight, this dosage corresponds in prevention to 1000 ppm of oxytetracycline in the feed, or an incorporation rate of 5 kg of medicated premix per ton of feed, and curative treatment at 2000 ppm of oxytetracycline in the feed, a incorporation rate of 10 kg of medicated premix per ton of feed.

Pigs: for daily feed intake of 25 g of feed per kg bodyweight, this dosage corresponds in prevention to 1000 ppm of oxytetracycline in the feed, or an incorporation rate of 5 kg medicated premix per ton of feed, and treatment to 2000 ppm oxytetracycline in the food, is an incorporation rate of 10 kg per medicated premix ton of feed.

INTERACTION WITH OTHER MEDICAMENTS

Divalent or trivalent cations (Mg, Fe, Al, Ca) can chelate tetracyclines. Tetracyclines should not be administered with antacids, gels based on aluminum, preparations based mineral as insoluble complexes are formed, which reduces the absorption of the antibiotic.

PREGNANCY AND LACTATION

The oxytetracycline showed no sign of embryotoxicity or teratogenicity in laboratory animals. In mammals, oxytetracycline through the placenta, causing tooth staining and slower fetal growth. The tetracyclines are found in breast milk. Product safety has not been evaluated in pregnant and lactating animals. Use of the product in pregnant and lactating animals will be an evaluation of the benefit / risk balance by the veterinarian.

OVERDOSE AND TREATMENT

Refer to section "SIDE EFFECT".

SIDE EFFECT

As for all tetracyclines, adverse effects were noted including gastrointestinal disturbances and less frequently allergic reactions and photosensitivity.

WITHDRAWAL TIME

Sheep, goats, pigs and Poultry (chickens, turkeys, ducks):

Meat and offal: 14 days

Egg: Zero days

SHELF LIFE: 36 months

SHELF LIFE after first opening & reconstitution: 24 hours

STORAGE

Store at temperature not exceeding 30°C. Strictly avoid light & heat exposure.

PACKAGING

The powder filled 25 kg in multi-wall bag.

DATE OF REVISION: 13 May 2019



Product registration Holder & Manufactured By:

SHENNONG ANIMAL HEALTH (M) SDN BHD

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,

Taman Perindustrian USJ 1,

47610 Subang Jaya, Selangor, Malaysia.