

Tetra-C 74% w/w Water Soluble Powder

Tetra-C 74% w/w Water Soluble Powder appears to be a yellow to brownish yellow color powder.

Composition:

Each g contains

Tetracycline hydrochloride 800 mg
(equivalent to Tetracycline 740 mg)

Pharmacodynamics:

The tetracyclines are bacteriostatic antibiotics that inhibit protein synthesis in susceptible microorganisms. After they diffuse through the outer cell membrane, an active carrier mediated process transports the drug through the inner cytoplasmic membrane. Once inside the cell, they bind irreversibly to receptors on the 30S subunit of the bacterial ribosome where they interfere with the binding of the aminoacyltransfer RNA to the acceptor site on the messenger RNA ribosome complex. This effectively prevents the addition of amino acids to the elongating peptide chain, inhibiting protein synthesis.

Pharmacokinetics:

The tetracycline group members vary in lipid solubility: it is this property which seems largely to determine their distribution and rate of excretion. They enter most tissues and body fluids with the exception of cerebrospinal fluid. As a result of chelation with calcium, tetracyclines become bound at active sites of ossification and in developing teeth. They cross the placenta to reach the foetus and are secreted in milk where they reach concentrations approximately those of serum. Most tetracyclines are excreted unchanged in urine and to a lesser extent, in bile. Tetracyclines undergo enterohepatic circulation, with much of the compound excreted in bile being reabsorbed from the intestine. This process contributes to the half-life of 6 - 10 hours, longer than might be anticipated with substances that are eliminated mainly by renal excretion.

Indications:

For use against diseases caused by *Clostridium perfringens* in chickens and those caused by *Actinobacillus* (*Haemophilus*) *pleuropneumoniae* in pigs.

Directions for Use:

For administration in the drinking water for up to 5 days. Allow birds and animals access to medicated water only during treatment.

Medicated water should be prepared daily and any medicated water that is not consumed within 24 hours should be discarded.

Chickens: Use 1 sachet (500 g) Tetra-C 74% to medicate 666 litres of water, providing 370 g of active ingredient per 666 litres. This is approximately equivalent to 60 mg tetracycline hydrochloride/kg bodyweight daily.

Pigs: Use 1 sachet (500 g) Tetra-C 74% to medicate 1110 litres of water, providing 370 g of active ingredient per 1110 litres. This is approximately equivalent to 40 mg tetracycline hydrochloride/kg bodyweight daily.

To allow ready dissolution of Tetra-C 74%, dissolve the required amount of product in a few litres of water before addition to the header tank. In areas where water is very hard it is advisable to add Tetra-C 74% at a rate of 1 sachet (500 g) to 370 litres of header tank water to ensure complete dissolution.

Mode of Administration:

Oral administration.

Contraindication:

None.

Warning and Precautions:

Special warnings for each target species

None known.

Special precautions for use

i) Special precautions for use in animals

None known.

ii) Special precautions to be taken by the person administering the medicinal product to animals

Wear suitable protective clothing, eye/face protection and impervious gloves before using the product.

Wash hands and exposed skin after handling the product.

Do not handle this product if you are allergic to tetracyclines.

In case of eye contact, rinse abundantly with water.

If you develop symptoms following exposure, such as skin rash, skin or eye irritation, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Interactions with Other Medicaments:

None known.

Statement on Usage During Pregnancy and Lactation:

No specific restrictions.

Adverse Effects / Undesirable Effects:

None known.

Overdose and Treatment:

None known.

Withdrawal Period (s):

Animals must not be slaughtered for human consumption during treatment. Animals may be slaughtered for human consumption after the following periods from the last treatment: Chickens: 3 days. Pigs: 6 days.

Do not use in laying birds producing eggs for human consumption.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure.

Keep medicine out of reach of children/

Jauhkan ubat dari kanak-kanak.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf-life as packaged for sale: 2 years

Shelf-life after first opening: 7 days

Shelf-life after reconstitution: 24 hours

Packaging Available:

500g & 1kg.

Date of revision:

01-03-2017

Product Holder/ Manufacturer:

NAM PHARMA SDN BHD

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833

