

SN-Paracetamol 500 mg/g

Water Soluble Powder
(FOR VETERINARY USE ONLY)

COMPOSITION

Paracetamol 500 mg
Exp q.s to 1 g

PRODUCT DESCRIPTION

The finished product is a white color powder.

TARGET SPECIES

Pigs (weaned pigs)

PHARMACODYNAMIC

Paracetamol or acetaminophen or N-acetyl-para-aminophenol is a derivative of para-aminophenol with analgesic and antipyretic properties. Its antipyretic effect is explained by its capacity to inhibit cerebral cyclo-oxygenases. Paracetamol is a weak inhibitor of COX-1 synthesis and therefore does not cause adverse effects on the gastrointestinal tract or changes in platelet aggregation.

PHARMACOKINETIC

Absorption: Oral absorption of paracetamol is rapid and almost completely (bioavailability of 90% after administration in drinking water). Maximum concentrations are reached in a little less than 2 hours after ingestion.

Metabolism: Paracetamol is mainly metabolised in the liver. The two major metabolic pathways are glucuronidation and sulfoconjugation. The latter pathway is rapidly saturable at dosages above therapeutic doses. A minor route, catalyzed by cytochrome P450, results in the formation of an intermediate reagent, N-acetyl-benzoquinoneimine which, under normal conditions of use, is rapidly detoxified by reduced glutathione and eliminated in urine after conjugation with cysteine and mercapturic acid. On the other hand, during mass poisoning, the quantity of this toxic metabolite is increased.

Elimination: Elimination is essentially urinary. In pigs, 63% of the ingested dose is eliminated by the kidney in 24 hours mainly in glucuronide-conjugated and sulfoconjugated form. Less than 5% is eliminated in unchanged form. The elimination half-life is about 4 hours. 63% of the ingested dose is eliminated by the kidney in 24 hours mainly in glucuronide-conjugated and sulfoconjugated form. Less than 5% is eliminated in unchanged form. The elimination half-life is about 4 hours.

INDICATIONS

In weaned pigs:

Symptomatic treatment of fever, in respiratory infection, if necessary in combination with appropriate anti-infectious treatment.

WARNINGS AND PRECAUTIONS

Special warnings for each target species:

Animals with low water consumption or poor general condition should be treated parenterally.

In the event of both viral and bacterial infection, appropriate anti-infective therapy should be administered concomitantly.

Special precautions for use:

i. Special precautions for use in animals:

A decrease of hyperthermia is expected 12-24 hours after initiation of treatment depending on the consumption of medicated water.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Wear suitable protective clothing, gloves and face / face protection. In case of contact with skin and eyes, rinse immediately with plenty of water. If symptoms persist, seek medical attention. To avoid any risk of ingestion, it is recommended not to eat or drink while using the product and to wash hands after handling the product. If swallowed, seek medical advice.

Do not handle the product if you are allergic to paracetamol.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

CONTRA-INDICATIONS

- Do not use in animals with known hypersensitivity to paracetamol or other constituents of the product,
- Do not use in animal with severe hepatic impairment,
- Do not use in animal with severe renal impairment. See also section 'INTERACTION WITH OTHER MEDICAMENTS'
- Do not use in animal with dehydration or hypovolemia.

ADMINISTRATION & DOSAGE

Oral use

30 mg of paracetamol per kg body weight per day, as long as the pigs are suffering from hyperthermia for a maximum of 5 days.

The product is administered continuously in the drinking water, at a dose of 0.6 g of product per 10 kg of live weight.

Depending on their clinical status, the animals consume a variable amount of water supplemented with medication. To ensure proper dosage, the concentration in the drinking water must be adjusted accordingly.

Recommendation for dissolution: pour the product in warm water preferably (30 to 35°C). Shake the preparation for 5 minutes to homogenize the drug solution. Then adjust the amount of water to obtain the required concentration and stir again until the solution is homogeneous.

INTERACTION WITH OTHER MEDICAMENTS

Concurrent administration of nephrotoxic drugs should be avoided.

PREGNANCY AND LACTATION

Studies in laboratory animals have not shown teratogenic or fetotoxic effects at the dose used in therapeutics.

Administration of the specialty to the pregnant and lactating sow, three times the therapeutic dose, did not result in any adverse effects.

OVERDOSE AND TREATMENT

After administration of 5 times the recommended dose of paracetamol, liquid faeces with solid particles may occasionally occur. It does not have any effect on general body condition of animals.

Acetylcysteine can be used in case of accidental overdose.

SIDE EFFECT

In rare cases, at therapeutic doses, transient soft faeces can occur and can persist up to 8 days after the withdrawal of administration. It does not have any effect on general condition of animals, and resolve without any specific treatment.

WITHDRAWAL TIME

Meat and offal: zero days.

SHELF LIFE: 36 months

SHELF LIFE after first opening & reconstitution: 24 hours

STORAGE

Store in room temperature ($\leq 30^{\circ}\text{C}$). Strictly avoid light & heat exposure.

PACKAGING

The powder filled 200 g or 500 g in aluminum sachets, by automatic filling machine and sealing, and packed in carton box.

DATE OF REVISION: 31 May 2019



Product registration holder & Manufactured by:

SHENNONG ANIMAL HEALTH (M) SDN BHD

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,

Taman Perindustrian USJ 1,

47610 Subang Jaya, Selangor, Malaysia.