

COTRIM Water Soluble Powder

Cotrim Water Soluble Powder appears to be a buff or light yellow color powder.

Composition:

Each g contains

Sulfachloropyridazine Sodium

625 mg

Trimethoprim

125 mg

Pharmacodynamics:

The antibacterial activity is based on a double sequential inhibition of microbial synthesis of tetrahydrofolic acid. Trimethoprim is an antibacterial derived from diaminopyrimidine having bactericidal and synergistic activity with sulfonamides. Sulfachloropyridazine is an antibacterial that works by preventing bacterial growth. The combination of the two active substances gives synergistic effect, which lower the doses of the active substances and enhance the bactericidal effect positively. It broadens the spectrum of activity and slow down the emergence of bacterial resistance.

Resistance develops slowly through transferable elements and results from a mutation chromosome as a result of one or more of the mechanisms described below:

- Permeability barrier and / or active efflux;
- Enzymes insensitive natural targets;
- Adjustable changes of the target enzyme;
- Recombination or mutational changes of the target enzyme;
- Acquired resistance by target enzymes resistant to the drug.

Pharmacokinetics:

Both components are rapidly absorbed after oral administration in fasted and fed animals and distributed in the body.

Indications:

Treatment of infections caused by bacteria susceptible to trimethoprim / sulfachloropyridazine, taking into account the ability of the antibiotic to reach the site of infection at the effective concentrations.

Recommended Dosage:

32mg Cotrim per kg body weight per day (= 24 mg of active substance in total) for 3-6 days.

Administration mode:

Chicken: Mix Cotrim in drinking water for 3-6 days.

Chicken	Age (weeks)	Dose: mg / liter of water
Broilers	1 - 4	160
	5 - 8	240
Breeders	1 - 8	160
	9 - 20	240
Hens	> 20	240

Preparation of drug drinking water: Half fill the drinking water container, add the prescribed amount of Cotrim, mix well and add the remaining water. The medicated drinking water is to be used within 24 hours of preparation.

Mode of Administration:

Oral administration.

Contraindication:

Do not use in laying hens producing eggs for human consumption. Do not use in hypersensitive animals or animals that have liver or kidney disease.

Warning and Precautions:

Special warnings for each target species

Avoid underdosing. Sick animals may drink less water than normal. If animals can be isolated, the concentration of Cotrim Water Soluble Powder can be increased proportionately in the water.

Special precautions for use in animals

The sensitivity of the target strains in relation to the product can vary. Susceptibility testing may be needed before treatment.

Water should always be present during the treatment period to avoid crystalluria.

Special precautions for the person administering the veterinary medicinal product to animals

Not applicable.

Interactions with Other Medicaments:

Do not use with other sulfonamides. There may be an interaction during concomitant treatment with an active substance that affects the liver metabolism and hematopoiesis.

Statement on Usage During Pregnancy and Lactation:

Do not use in laying hens.

Side effects:

Sulfonamides can affect renal function (crystalluria, hematuria, urinary obstruction) and hematopoiesis (thrombocytopenia, anemia).

Overdose and Treatment:

No adverse effect has been recorded.

Withdrawal Period (s):

Chicken:

Meat: 7 days.

Eggs: Not for use in chickens laying eggs intended for human consumption.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure.

Keep out of reach of children/Jauhkan dari kanak-kanak.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf life after first opening: 7 days.

Shelf life after reconstitution: 24 hours.

Packaging Available:

73 g & 100 g

Date of revision:

23-03-2017

Product Holder/ Manufacturer:

NAM PHARMA SDN. BHD. (768370-H)

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833