

PACKAGE INSERT

1. Brand Name: AMOX EMV 500 mg/g Water Soluble Powder

2. Dosage form: Oral solid - powder to be dissolved in water

Name of strength of active substance and excipient

Amoxicillin Trihydrate.....500 mg

Dextrose anhydrous q.s to.....1 g

3. Product description: white to yellowish white powder with characteristic odor

4. Indication:

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin.

Not effective against beta-lactamase producing organisms.

Use of the product should be based on susceptibility testing and it should take into account official and local antimicrobial policies.

Pigs: For the treatment of salmonellosis and pasteurellosis.

5. Pharmacodynamics:

Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against Gram positive and Gram-negative bacteria. It owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

6. Pharmacokinetics:

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids.

Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

7. Environmental Properties: Not applicable

Target species: Chickens, ducks, turkeys, pigs.

8. Recommended dose:

8.1 Chickens, turkeys and ducks

The product is administered in the drinking water. Prepare the solution with fresh tap water immediately before use. Once opened, use the contents of one sachet immediately. Any unused medicated water should be discarded after 12 hours.

In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated.

The following formula may be used to calculate the amount of product required per day (in grams):

$$\frac{\text{Number of birds} \times \text{average live weight (kg)}}{25 \text{ (for 20 mg/kg) or } 33 \text{ (for 15 mg/kg)}}$$

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing. The uptake of medicated water depends on the clinical condition of the birds. In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted taking into account water intake.

Chickens:

The recommended dosage is 15 mg amoxicillin trihydrate (30 mg AMOX EMV 500 mg/g Water Soluble Powder) per kg body weight. The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks:

Recommended dosage is 20 mg amoxicillin trihydrate (40 mg AMOX EMV 500 mg/g Water Soluble Powder) per kg bodyweight for 3 consecutive days.

Turkeys:

Recommended dosage is 15-20 mg amoxicillin trihydrate (30-40 mg AMOX EMV 500 mg/g Water Soluble Powder) per kg bodyweight for 3 days or in severe cases for 5 days

8.2 Pigs:

For the medication of Pigs, this product may be administered in the drinking water or administered by addition to liquid feeds produced with commercial feed. It may not be used in dry feeds.

8.2.1. Administration via the drinking water

a. Administer in the drinking water to give 20mg/kg bodyweight daily. The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days. Dissolve the contents of one sachet (100g of the product) in the requisite quantity of water immediately before use.

b. For administration via the drinking water, the solution should be prepared with fresh potable water.

- i. Any medicated water which is not consumed within 12 hours should be discarded.
- ii. In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst on treatment

8.2.2. Administration in liquid feed

a. Administer in the liquid feed, to give 20mg amoxicillin trihydrate/kg bodyweight daily for up to 5 days. Medicated feed should be freshly prepared on at least 3 occasions per day over the treatment period. The daily dose should be calculated based on the number of animals and average weight and then divided by the number of feed lots prepared in the day.

b. Medicated liquid feed should be prepared with fresh potable water.

c. Dissolve the required amount of this product in water at a rate of 25g/L before addition to the feed. This may be done by dissolving the contents of one sachet (100g of the product) in approximately 4L of water immediately before use.

d. After adding the product to some or all of the water needed to make the liquid feed, ensure the product is fully dissolved. Dissolution of the product can take up to 10 minutes. This medicated water can then be mixed with the dry complete meal and if appropriate, the remaining water. The system used should ensure that the medicated water is evenly distributed into the feed. Once prepared the final medicated liquid feed should be fed to the pigs immediately.

e. The medicated liquid feed should not be fermented and should not be stored.

f. Stability of amoxicillin in all commercial feeds has not been established. In order to ensure that any loss of amoxicillin activity is minimized, the quantity of medicated liquid feed prepared should not exceed the amount of feed which will be consumed within 4 hours.

- i. Any medicated liquid feed which is not consumed within 4 hours should be discarded
- ii. Although restricted access to other water supplies would help ensure medicated liquid feed is consumed, separate clean potable water should remain available at all times for welfare reasons.

9. Contraindication:

Do not be administered to rabbits, hamsters, gerbils and guinea pigs, or to birds producing eggs intended for human consumption.

10. Special warning and precautions:

Special warnings for each target species

None

Special precautions for use

Special precautions for use in animals

Pigs: The uptake of medication by animals can be altered as a consequence of illness. In case of insufficient uptake of feed or water, animals should be treated parenterally.

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa.

Allergic reactions to these substances may occasionally be serious.

- 1) Do not handle this product if you know you are sensitive or if you have been advised not to work with such preparations.
- 2) Handle this product with great care to avoid exposure, taking all recommended precautions.
- 3) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Avoid inhalation of dust. Wear either a disposable half-mask respirator conforming to European Standard EN149 or a non-disposal respirator to European Standard EN140 with a filter to EN143

Wear gloves during preparation and administration of medicated water or liquid feed

Wash any exposed skin after handling the product or medicated water or feed.

Wash hands after use

11. **Side effects:**

None known

12. **Interaction of other medicaments:**

None known.

13. **Pregnancy and lactation:**

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.

Use only according to the benefit/risk assessment of the responsible veterinarian.

14. **Symptoms and Treatment of overdose:**

No problems with overdosage have been reported. Treatment should be symptomatic and no specific antidote is available.

15. **Storage condition:**

Do not store above 30°C.
Store at dry place

16. **Shelf life:**

The product packaged for sale has a shelf-life of 2 years.

The shelf-life of the first-opening container: 24 hours

Shelf-life after dilution according to directions: 12 hours.

17. **Withdrawal Period:**

Chickens (meat & offal): 1 day

Ducks (meat & offal): 9 days

Turkeys (meat & offal): 5 days

Pigs (meat & offal): 2 days

Not for use in laying birds producing eggs for human consumption.

18. **Manufacturer:**

Emivest Feedmill Vietnam Company Limited

Address: No. A-11C-CN Bau Bang Industrial Park, Lai uyen Town, Bau Bang District, Binh Duong Province, Vietnam

19. **Importer and Distributor:**

F.E VENTURE SDN BHD

Lot 3 & 5, Jalan PJS 11/8, Bandar Sunway, 46150 Petaling Jaya, Selangor Darul Ehsan, Malaysia