

TRIMETOSOL 80/400 G/KG POWDER

Description and Composition

TRIMETOSOL 80/400 G/KG POWDER is a white to off white powder. Each kg contains 80g Trimethoprim and 400g Sulfamethoxazole as its active ingredients.

Pharmacodynamics

Trimethoprim and sulfamethoxazole have a broad spectrum of activity against gram-positive and gram-negative bacteria including *Streptococcus* spp., *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, *Haemophilus parasuis*, *Avibacterium paragallinarum* and *E. coli* in vitro. Sulphonamides block the conversion of para-aminobenzoic acid to dihydrofolic acid. Its effect is bacteriostatic.

Trimethoprim inhibits dihydrofolic acid reductase, which converts dihydrofolic into tetrahydrofolic acid. The effect of trimethoprim is bacteriostatic and in combination with sulphonamides it is bactericidal. Sulphonamides and trimethoprim thus cause a successive blockage of two enzymes that play an important role in the metabolism of bacteria and protozoa. Their effect is synergistic.

Bacterial resistance to trimethoprim and to sulphonamides can be mediated via 5 main mechanisms: (1) changes in the permeability barrier and/or efflux pumps, (2) naturally insensitive target enzymes, (3) changes in the target enzymes, (4) mutational or recombinational changes in the target enzymes, and (5) acquired resistance by drug-resistant target enzymes.

Pharmacokinetics

Following oral administration, trimethoprim and sulfamethoxazole are rapidly and almost completely absorbed from the gut. The bioavailability of sulfamethoxazole is slightly higher than that of trimethoprim. It is distributed to all tissues except the brain. The highest concentrations can be found in the lungs, the liver and the kidneys.

Sulphonamides are metabolised in various ways. The degree of acetylation, hydroxylation and glucuronidation depends, among other things, on the species and age of the animal. Trimethoprim is metabolised to a large extent in the liver. Important metabolic pathways are O-methylation, N-oxidation in the ring structure and alpha hydroxylation. Sulfamethoxazole and trimethoprim are primarily excreted through the kidneys.

Indication

Treatment of bacterial diseases and diarrhoea caused by *E. coli* susceptible to trimethoprim and sulfamethoxazole in piglets and chickens.

Recommended Dose

The following dose is mixed with feed and administered orally once a day for 5 to 7 days. 551g of product per tonne of feed.

Route of Administration

For oral administration via feed.

Contraindication

Do not use in animals suffering from severe liver or kidney disease, oliguria or anuria.

Do not use in animals with impaired haematopoietic systems.

Do not use in case of known hypersensitivity to sulphonamides or trimethoprim or any of the excipients.

Warning and Precautions

Special warnings for each target species:

- Do not administer to pigs over 4 months of age.
- Do not use in laying hens.

Special precautions for use:**Special precautions for use in animals:**

- Do not administer to animals showing shock and hypersensitivity reactions to this drug and other sulfa antibiotics.
- Use according to the instructions and prescription of a veterinarian.
- Do not use it as safety and effectiveness have not been established except for the mentioned target species.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- The handler should wear protective equipment such as gloves, mask, and protective clothing to avoid contact or inhalation.
- In case of contact with the skin or eyes when handling the drug, wash it off immediately. Seek medical attention immediately if abnormal symptoms appear.

Interaction with Other Medicaments

Do not combine with other veterinary medicinal products.

Pregnancy and Lactation

Do not use in laying hens.

Side Effects

- Sulfa drugs can cause disorders of the urinary system, such as crystal urine, hematuria, and urinary tract obstruction.
- Sulfa drugs can cause hypersensitivity reactions and skin rashes.
- Long-term administration of sulfa and trimethoprim may cause anemia, leukopenia, thrombocytopenia, etc.

Symptoms and Treatment of Overdose

Chronic overdose in chickens will result in a strongly diminished water- and feed intake and retarded growth.

Storage Condition

Store below 30°C.

Shelf life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Periods

Pig: 63 days

Chickens: 14 days

Do not use in laying hens.

Packing

10kg and 25kg.

Revised date: 18.12.2020

	Manufacturer & Product Registration Holder Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com
---	---