

PACIFIC CTC 15% w/w Granule

Active substance & strength: Each 1 kg contains: 150 grams of Chlortetracycline Hydrochloride

Product Description: Brown to dark brown granule, not lumpy or moldy, no unpleasant odor.

Pharmacodynamics: Chlortetracycline is a natural broad-spectrum antibiotic of the tetracycline group. Its activity is bacteriostatic by interfering with the protein synthesis at the S30 ribosome sub-unit. At recommended dosages it has no pharmacological effects on cardiovascular, nervous or other body systems.

Pharmacokinetics: When dosed orally it is quickly absorbed into the blood stream. In-feed dosing provides steady to near steady state plasma levels. This provides effective concentrations in various tissues, including lungs and other respiratory tissues. The major metabolite is 4-epi-chlortetracycline.

Route of Administration: Oral

Recommended Dose:

Chickens:

- 1) Control of infectious synovitis caused by *Mycoplasma synoviae* susceptible to chlortetracycline, 110-220 grams per tonne of feed continuously for 7 to 14 days.
- 2) Control of chronic respiratory disease and air sac infection caused by *Mycoplasma gallisepticum* and *Escherichia coli* susceptible to chlortetracycline, 220-440 grams per tonne of feed continuously for 7 to 14 days.
- 3) Reduction of mortality due to *Escherichia coli* infections susceptible to chlortetracycline, 550 grams per tonne of feed, for 5 days.

Swine:

- 1) Control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis* susceptible to chlortetracycline, 22 milligrams per kg of body weight per day, not more than 14 days.
- 2) Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline, 22 milligrams per kg of body weight per day. Feed approximately 440 g/tonne, varying with body weight and feed consumption to provide 22 mg/kg per day, not more than 14 days.
- 3) Reduction in the incidence of cervical lymphadenitis (jowl abscesses) caused by group E *Streptococci* susceptible to chlortetracycline, 55-110 grams per tonne of feed.

Swine (Breeding): Control of leptospirosis (reducing the instances of abortion and shedding of *Leptospirae*) caused by *Leptospira pomona* susceptible to chlortetracycline, 440 grams per tonne of feed continuously for not more than 14 days.

Withdrawal Period(s): No withdrawal period is required

Warnings & Precautions: Special precautions for use in animals: Official, national and regional antimicrobial policies should be taken into account when the product is used. Special precautions to be taken by the person administering the veterinary medicinal product to animals. If you know you are hypersensitive (allergic) to chlortetracycline, do not handle the product. When handling this product, inhalation of the dust must be avoided by wearing a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143. Avoid contact with skin and eyes. Gloves should be worn whilst handling this product. If contact with skin or eyes occurs, wash area immediately with clean fresh water. If irritation persists seek medical attention. Do not eat, drink or smoke whilst handling the product. Hands and exposed skin should be washed thoroughly after use.

Contraindications: Do not use in ruminants

Pregnancy and Lactation: No problems known

Mode of Administration: Orally into feed

Interactions with Other Medicaments: Chlortetracycline is sometimes used in association with other antimicrobials, notably sulphadimidine, penicillin and neomycin. There are no known adverse reactions with these or any other substances.

Adverse Effects/Undesirable Effects: The product is of low toxicity and side effects are rarely encountered. If suspected adverse reactions occur, treatment should be discontinued immediately.

Overdose and Treatment: Chlortetracycline is of low toxicity and there is a wide safety margin at the recommended dosage. On rare occasions overdosage may cause diarrhoea and over growth of yeast and fungi. Under such conditions, withdraw medication and apply appropriate treatment.

Storage condition: Store in cool (<25 °C) and dry place without sunlight.

Packing: 25kg bag

Manufacturer : Jinhe Biotechnology Co. Ltd. No. 71 Xinping Road, Tuoketuo County, Hohhot 010200, Inner Mongolia, China

Marketing authorization holder: QL Pacific Vet Group Sdn Bhd. No 886-C8, Jalan Subang 9, Taman Perindustrian Subang, 47600 Subang Jaya Selangor, Malaysia

Date of Revision of Package Insert: 01 Mar 2021