

Pet-Amox 62.5 mg Tablet

Amoxicillin 50 mg and Clavulanic acid 12.5 mg tablets

COMPOSITION:

One tablet contains:
Amoxicillin (as amoxicillin trihydrate) 50 mg
Clavulanic Acid (as potassium salt) 12.5 mg

DESCRIPTION

The finished product is beige, oval shaped, scored tablets.

TARGET SPECIES

Dogs and cats

PHARMACODYNAMICS

Amoxicillin is an aminobenzylpenicillin from the β -lactam penicillin family which prevents the bacterial cell wall formation by interfering with the final step of peptidoglycan synthesis.

Clavulanic acid is an irreversible inhibitor of intracellular and extracellular β -lactamases which protects amoxicillin from inactivation by many β -lactamases.

Amoxicillin/ clavulanate has a wide range of activity which includes β -lactamase producing strains of both Gram-positive and Gram-negative aerobes, facultative anaerobes and obligate anaerobes.

PHARMACOKINETICS

After oral administration at the recommended dose in dogs and cats, the absorption of amoxicillin and clavulanic acid is fast. In dogs, the maximum plasma concentration of amoxicillin of 8.5 $\mu\text{g/ml}$ is reached in 1.4 hours and the maximum plasma concentration of clavulanic acid of 0.9 $\mu\text{g/ml}$ is reached in 0.9 hours. Half life is 1 hour in dogs for both substances.

In cats, the maximum plasma concentration of amoxicillin of 6.6 $\mu\text{g/ml}$ is reached in 1.8 hours and the maximum plasma concentration of clavulanic acid of 3.7 $\mu\text{g/ml}$ is reached in 0.75 hours. Half life is 1 to 2 hours in cats for both substances.

Elimination is also fast. 12 % of the amoxicillin and 17 % of clavulanic acid is excreted in the urine. The remainder is excreted as inactive metabolites.

After repeated oral administration of the recommended dose in dogs and cats, there is no accumulation of amoxicillin or clavulanic acid and the steady state is reached rapidly after first administration.

INDICATION

In dogs: treatment or adjunctive treatment of periodontal infections caused by bacteria susceptible to amoxicillin in

combination with clavulanic acid i.e. *Pasteurella* spp, *Streptococcus* spp and *Escherichia coli*.

In cats: treatment of skin infections (including wounds and abscesses) caused by bacteria susceptible to amoxicillin in combination with clavulanic acid i.e. *Pasteurella* spp, *Staphylococcus* spp, *Streptococcus* spp and *Escherichia coli*.

ROUTE OF ADMINISTRATION

For oral administration

RECOMMENDED DOSE

To ensure the correct dosage, body weight should be determined as accurately as possible to avoid under-dosing. The recommended dose of the product is 10 mg amoxicillin / 2.5 mg clavulanic acid per kg body weight twice a day by the oral route in dogs and cats, i.e. 1 tablet per 5 kg body weight every 12 h, according to the following table:

Bodyweight (kg)	Number of tablets twice daily
[1.0- 2.5]	$\frac{1}{2}$
[2.6- 5.0]	1
[5.1- 7.5]	1 $\frac{1}{2}$
[7.6- 10.0]	2

In severe infections in each target species, the dose may be doubled to 20 mg amoxicillin/5 mg clavulanic acid/kg body weight twice daily.

Duration of treatment:

- 7 days for the treatment of periodontal infections in dogs.

- 7 days for the treatment of skin infections in cats (including wounds and abscesses). The clinical status of animals should be re-evaluated after 7 days and the treatment prolonged for a further 7 days if necessary. Severe cases of skin infection may require an even longer duration of treatment and this should be at the discretion of the responsible veterinarian.

CONTRAINDICATIONS

Do not use in cases of hypersensitivity to penicillins or other substances of the β -lactam group or to any of the excipients. Do not administer to gerbils, guinea pigs, hamsters, rabbits and chinchillas.

Do not administer to horses and ruminating animals.

Do not use in animals with serious dysfunction of the kidneys accompanied by anuria or oliguria.

Do not use in cases of known resistance to the combination of amoxicillin and clavulanic acid.

INTERACTION WITH OTHER MEDICAL PRODUCT

The bactericidal activity of amoxicillin may be reduced by the simultaneous use of bacteriostatic substances such as macrolides, tetracyclines, sulfonamides and chloramphenicol.

The potential for allergic cross-reactivity with other penicillins should be considered.

Penicillins may increase the effect of aminoglycosides.

WARNINGS AND PRECAUTIONS

Special warnings for each target species

None

Special precautions for use

Special precautions for use in animals

In animals with impaired liver and kidney function, the use of the product should be subject to a risk/benefit evaluation by the veterinary surgeon and the posology evaluated carefully.

Caution is advised in the use in small herbivores other than those in Contraindications.

Use of the product should be based on susceptibility testing. Use of the product deviating from the instructions may increase the prevalence of bacteria resistant to amoxicillin/ clavulanic acid may decrease the effectiveness of treatment with other β -lactam antibiotics, due to the potential for cross resistance. Use of the product should take into account official and local antimicrobial policies. Do not use in cases of bacteria sensitive to narrow spectrum penicillins or to amoxicillin as a single substance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and *vice versa*. Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.

2. Handle this product with great care to avoid exposure, taking all recommended precautions.

3. If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the doctor this warning.

Swelling of the face, lips or eyes or difficulty breathing are more serious symptoms and require urgent medical attention.

Wash hands after handling the tablets.

- Keep out of reach of children
- Jauhkan daripada kanak-kanak

PREGNANCY AND LACTATION

The safety of the product has not been established during pregnancy and lactation. Laboratory studies in rats have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. Use only in accordance with the benefit/ risk assessment by the responsible veterinarian.

SIDE EFFECTS

Vomiting and diarrhoea may be observed very rarely. Treatment may be discontinued depending on the severity of the undesirable effects and a benefit/risk evaluation by the veterinary surgeon.

Hypersensitivity reactions (allergic skin reactions, anaphylaxis) may be observed very rarely. In these cases, administration should be discontinued and a symptomatic treatment given.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

SYMPTOMS AND TREATMENT OF OVERDOSE

At three times the recommended dose for a period of 28 days, a decrease in cholesterol values and episodes of vomiting were observed in cats and diarrhoea was observed in dogs. In the event of an overdose symptomatic treatment is advised.

STORAGE CONDITION

Store in a dry place not exceeding 30°C. Protect from light.

SHELF LIFE: 2 years

SHELF LIFE AFTER OPENING: 24 hours

WITHDRAWAL PERIOD: Not applicable

PACKAGING

Each box contains 100 tablets in 10 blister strips.

REVISED DATE: Dec 14, 2020



Manufactured by:
Vemedim Veterinary Drugs And Probiotics Co., Ltd
(Vemedim Animal Health)
Song Hau Industrial Zone, Chau Thanh District,
Hau Giang Province, Vietnam

Product Registration Holder:
Vemedim Sdn Bhd
15, Jalan USJ 1/31, 47600 Subang Jaya
Selangor, Darul Ehsan, Malaysia