

TRIMETOSOL 80/400 MG/ML SUSPENSION

Description and Composition

TRIMETOSOL 80/400 MG/ML SUSPENSION is a white to light brown suspension. Each ml contains 80mg Trimethoprim and 400mg Sulfamethoxazole as its active ingredients.

Pharmacodynamics

Trimethoprim and sulfamethoxazole have a broad spectrum of activity against gram-positive and gram-negative bacteria including *Streptococcus* spp., *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, *Haemophilus parasuis*, *Avibacterium paragallinarum* and *E. coli* in vitro. Sulphonamides block the conversion of para-aminobenzoic acid to dihydrofolic acid. Its effect is bacteriostatic.

Trimethoprim inhibits dihydrofolic acid reductase, which converts dihydrofolic into tetrahydrofolic acid. The effect of trimethoprim is bacteriostatic and in combination with sulphonamides it is bactericidal. Sulphonamides and trimethoprim thus cause a successive blockage of two enzymes that play an important role in the metabolism of bacteria and protozoa. Their effect is synergistic.

Bacterial resistance to trimethoprim and to sulphonamides can be mediated via 5 main mechanisms: (1) changes in the permeability barrier and/or efflux pumps, (2) naturally insensitive target enzymes, (3) changes in the target enzymes, (4) mutational or recombinational changes in the target enzymes, and (5) acquired resistance by drug-resistant target enzymes.

Pharmacokinetics

Following oral administration, trimethoprim and sulfamethoxazole are rapidly and almost completely absorbed from the gut. The bioavailability of sulfamethoxazole is slightly higher than that of trimethoprim. It is distributed to all tissues except the brain. The highest concentrations can be found in the lungs, the liver and the kidneys.

Sulphonamides are metabolised in various ways. The degree of acetylation, hydroxylation and glucuronidation depends, among other things, on the species and age of the animal. Trimethoprim is metabolised to a large extent in the liver. Important metabolic pathways are O-methylation, N-oxidation in the ring structure and alpha hydroxylation. Sulfamethoxazole and trimethoprim are primarily excreted through the kidneys.

Indication

Treatment of bacterial diseases and diarrhoea caused by *E. coli* susceptible to sulfamethoxazole and trimethoprim in piglets.

Recommended Dose

Piglets: 0.125ml of product per 2kg of body weight is administered orally once a day, for 3 to 5 days.

Route of Administration

By oral administration only. Shake well before use.

Contraindication

- Do not use this product in combination with methotrexate, warfarin, phenylbutazone, thiazide diuretics, salicylates, probenecid, phenytoin, etc.
- Do not administer with procaine, tetracycline, and methenamine as severe ionophore poisoning symptoms may appear.
- Do not take with antacids.

Warning and Precautions

Special warnings for each target species:

None known.

Special precautions for use**Special precautions for use in animals:**

- Do not administer to animals showing shock and hypersensitivity reactions to this drug and other sulfa antibiotics.
- Use as directed by veterinarian.
- Do not use for other target species besides mentioned in indication.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- The handler should wear protective equipment such as gloves, mask, and protective clothing. Avoid contact or inhalation.
- In case of contact with the skin or eyes when handling the drug, wash it off immediately. Seek medical attention.

Interaction with Other Medicaments

Do not combine with other veterinary medicinal products.

Pregnancy and Lactation

The safety of the veterinary medicinal product during pregnancy, lactation or lay has not been established.

Side Effects

- Sulfa drugs can cause urinary disorders such as crystal urine, hematuria, and urinary tract obstruction.
- Sulfa drugs may cause skin rash due to hypersensitivity reactions.
- Long-term administration of sulfa and trimethoprim causes anemia, leukopenia, and thrombocytopenia.

Symptoms and Treatment of Overdose

In pigs, an overdose of 2 ½ times the recommended dosage does not induce any adverse effects.

Storage Condition

Store below 30°C.

Shelf life

3 years.

After first opening the container, any product remaining after 24 hours should be discarded.

Withdrawal Period

63 days.

Packing

200ml, 500ml, 1L.

Revised date: 23.4.2021

	Manufacturer & Product Registration Holder Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com
---	---