

1. Name of the medical product

COMOZOL cream 2% w/w

2. Qualitative and Quantitative composition

Each gram contains:

Ketoconazole..... 20mg

3. Pharmaceutical form

White and homogeneous cream

4. Clinical particulars**4.1 Indications**

Comozol Cream 2% is indicated for topical application in the treatment of dermatophyte infections of the skin: tinea corporis, tinea cruris, tinea manus, and tinea pedis due to *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Microsporum canis*, and *Epidermophyton floccosum*, as well as in the treatment of cutaneous candidosis and tinea (pityriasis) versicolor.

- Comozol Cream 2% is also indicated for the treatment of seborrhoeic dermatitis, a skin condition associated with the presence of *Malassezia furfur*.

4.2 Posology and method of administration

- Cutaneous candidosis, tinea corporis, tinea cruris, tinea manus, tinea pedis and tinea (pityriasis) versicolor: it is recommended that Comozol Cream 2% be applied once daily to cover the affected and immediate surrounding area.

- Seborrhoeic dermatitis: Comozol Cream 2% should be applied to the affected area once or twice daily.

- The usual duration of treatment is: tinea versicolor 2 - 3 weeks, yeast infections 2 - 3 weeks, tinea cruris 2 - 4 weeks, tinea corporis 3 - 4 weeks, tinea pedis 4 - 6 weeks.

- The usual initial duration of treatment in seborrhoeic dermatitis is 2 to 4 weeks. Maintenance therapy is applied once or twice weekly in seborrhoeic dermatitis.

- Treatment should be continued, until a few days after disappearance of all symptoms. The diagnosis should be reconsidered if no clinical improvement is noted after 4 weeks of treatment.

Special populations

Paediatrics

There are limited data on the use of ketoconazole 2% cream in paediatric patients.

Administration

Topical administration to the skin.

For External Use Only.

4.3 Contraindications

COMOZOL Cream is contra-indicated in patients with a known hypersensitivity to any of the ingredients or to ketoconazole itself.

4.4 Special warnings and precautions for use

COMOZOL Cream is not for ophthalmic use.

If coadministered with a topical corticosteroid, to prevent a rebound effect after stopping a prolonged treatment with topical corticosteroids it is recommended to continue applying a mild topical corticosteroid in the morning and to apply COMOZOL Cream in the evening,

and to subsequently and gradually withdraw the topical corticosteroid therapy over a period of 2-3 weeks.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed.

4.6 Pregnancy and lactation

There are no adequate and well-controlled studies in pregnant or lactating women. Plasma concentrations of ketoconazole are not detectable after topical application of ketoconazole to the skin of non-pregnant humans. There are no known risks associated with the use of topical ketoconazole in pregnancy or lactation.

4.7 Effects on ability to drive and use machines

COMOZOL Cream has no influence on the ability to drive and use machines.

4.8 Undesirable effects

System Organ Class	Adverse Reactions		
	Category		
	Common	Uncommon	Not Known
Immune System Disorders		Hypersensitivity	
Skin and Subcutaneous Tissue Disorders	Skin burning sensation	Bullous eruption Dermatitis contact Rash Skin exfoliation Sticky skin	Urticaria
General Disorders and Administration Site Conditions	Application site erythema Application site pruritus	Application site bleeding Application site discomfort Application site dryness Application site inflammation Application site irritation Application site paresthesia Application site reaction	

4.9 Overdose

Topical Application

Excessive topical application may lead to erythema, oedema and a burning sensation, which will disappear upon discontinuation of the treatment.

Ingestion

In the event of accidental ingestion, supportive and symptomatic measures should be carried out.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antifungals for Topical Use, Imidazole and triazole derivatives

ATC Code: D01AC08

Ketoconazole inhibits the biosynthesis of ergosterol in fungi and changes the composition of other lipid components in the membrane.

Usually ketoconazole cream acts rapidly on pruritus, which is commonly seen in dermatophyte and yeast infections, as well as skin conditions associated with the presence of *Malassezia* spp. This symptomatic improvement is observed before the first signs of healing are observed.

Ketoconazole, a synthetic imidazole dioxolane derivative, has a potent antimycotic activity against dermatophytes such as *Trichophyton* spp., *Epidermophyton floccosum* and *Microsporum* spp. and against yeasts, including *Malassezia* spp. and *Candida* spp. The effect on *Malassezia* spp. is particularly pronounced.

5.2 Pharmacokinetic properties

Plasma concentrations of ketoconazole were not detectable after topical administration of Ketoconazole Cream in adults on the skin.

6. Pharmaceutical particulars

6.1 List of excipients

Propylene glycol
Stearyl alcohol
Cetanol
Sorbitanmonostearate
Isopropyl myristate
Polysorbate 60
Butylatedhydroxytoluene
Purified water

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store in well closed containers, below 30°C.

6.5 Nature and contents of container

10g/Tube/Box

7. Manufacturer

DAE HWA PHARMACEUTICAL CO., LTD.
495, Hanu-ro, Hoengseong-Eup, Hoengseong-Gun, Gangwon-Do, Republic of Korea

8. Product Registration Holder

Eucogen Sdn. Bhd.
No 6A, Jalan Sungai Burung U32/U,
Bukit Rimau, 40460 Shah Alam, Selangor, Malaysia.

9. Date of Revision of the text

Dec. 2020