



Diaziprim

Oral solution • Sulfadiazine, trimethoprim

Composition 1 ml contain

Active substance: Sulfadiazine 83.35 mg •
Trimethoprim 16.65 g.

Indications

Treatment of infections caused by microorganisms sensitive to sulfadiazine and trimethoprim. In calves, lambs, pigs, rabbits and poultry: prevention and treatment of respiratory and gastrointestinal infections.

Method and routes of administration

See the package insert before use.

Special warnings | Disposal of containers

See the package insert before use.

Withdrawal periods

Meat and offal: 12 days

Eggs: do not use in laying birds producing eggs for human consumption, 4 weeks before laying starts and during laying.

Storage conditions

Keep below 25 °C. Shelf life after first opening : 3 months. Shelf life after dilution in drinking water: 12 hours. Shelf life after dilution in milk: 2 hours.

Keep out of the sight and reach of children | Jauhkan daripada kanak-kanak
Controlled medicine | Ubat terkawal • For animal treatment only

Lot :

Mfg :

Exp :



HUVEPHARMA®

MAL

10 L

Manufactured by: **HUVEPHARMA SA** • 34, rue Jean Monnet • Z.I. d'Étriché • Segré • 49500 Segré-en-Anjou Bleu • France • tél. : +33 (0)2 41 92 11 11 • info.france@huvepharma.com

Registration holder: **RHONE MA MALAYSIA SDN BHD** (200001023279) • Lot 18A & 18B • Jalan 241 • Seksyen 51A • 46100 Petaling Jaya • Selangor • Malaysia

ORAL SOLUTION

CALVES • LAMBS • PIGS • POULTRY • RABBITS



Diaziprim Oral solution

Package leaflet for:

Composition

Active substance : Sulfadiazine 83.35 mg •
Trimethoprim 16.65 mg.

Pale yellow to ochre yellow clear solution.

Pharmacodynamics

Pharmacokinetics

important. The distribution is good in most of tissues and organs. Trimethoprim is rapidly absorbed after oral administration. It is widely distributed in the body. The two active ingredients are partially metabolized in the liver. Their elimination is essentially renal.

Indication

Recommended dosage and mode of administration

Calves and lambs: 12.5 mg of sulfadiazine and 2.5 mg of trimethoprim/kg body weight/12 hours, for 4-7 days, i.e. 1.5 ml of solution/10 kg body weight/12 hours, for 4-7 days, to mix with the milk feed (when adding the water).

Pigs, rabbits and poultry: 25 mg of sulfadiazine and 5 mg of trimethoprim/kg body weight/day for 4-7 days, i.e. 3 ml of solution-10 kg body weight/day for 4 to 7 days, to be diluted in drinking water or liquid food.

The uptake of medicated water or milk depends on the clinical condition of the animals. In order to obtain the correct dosage the concentration of sulfadiazine and trimethoprim has to be adjusted accordingly. The medicated water is only usable for 12 hours and should be made freshly every day. The medicated milk is only

usable for 2 hours and should be made freshly every day.

Contraindications

Do not use in case of hypersensitivity to sulfadiazine or trimethoprim. Do not use in animals with severe renal or hepatic dysfunction.

Special warnings for each target species

Do not administer to laying hens uses to produce eggs for human consumption.

Special precautions for use

Special precautions for use in animals: None known. Special precautions to be taken by the person administering the medicinal product to animals: People with known hypersensitivity to sulfadiazines should avoid contact with the product. Do not eat, drink or smoke while handling the product. If any symptom should appear, such as a cutaneous eruption, seek prompt medical advice and show the package leaflet or the label to the doctor. Swelling of the face, lips or eyes or respiratory difficulties are the most serious signs with require urgent medical attention. Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products, if appropriate: Any unused veterinary medicinal product or waste materials

derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Interaction with other medicaments

None known.

Use during pregnancy and lactation

Laboratory studies in animals have shown evidence of teratogenic and foetotoxic effects at doses exceeding the recommended therapeutic doses. The use is not recommended during pregnancy and lactation

Adverse effects

None known.

Overdose and treatment

None known.

Withdrawal period

Meat and offal: 12 days.

Eggs: do not use in laying birds producing eggs for human consumption, 4 weeks before laying starts and during laying.

Storage conditions

Keep below 25 °C. Shelf life after first opening : 3 months. Shelf life after dilution in drinking water: 12 hours. Shelf life after dilution in milk: 2 hours.

Packing: 2 L, 5 L and 10 L.

Keep out of the sight and reach of children | Jauhkan daripada kanak-kanak
Controlled medicine | Ubat terkawal • For animal treatment only



HUVEPHARMA

Manufactured by: **HUVEPHARMA SA** • 34, rue Jean Monnet • Z.I. d'Étriché • Segré • 49500 Segré-en-Anjou Bleu • France • tél. : +33 (0)2 41 92 11 11 • info.france@huvepharma.com

Registration holder: **RHONE MA MALAYSIA SDN BHD** (200001023279) • Lot 18A & 18B • Jalan 241 • Seksyen 51A • 46100 Petaling Jaya • Selangor • Malaysia