

OMS TMS 480 PREMIX POWDER

Composition

Each gm contains:

Sulfamethoxazole.....400.0 mg

Trimethoprim.....80.0 mg

Product Description

White or almost white powder

Pharmacodynamics

Trimethoprim and sulfamethoxazole have a broad spectrum of activity against gram-positive and gram-negative bacteria including *Streptococcus* spp. *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, *Haemophilus parasuis*, *Avibacterium paragallinarum* and *E. coli* in vitro. Sulphonamides block the conversion of para-aminobenzoic acid to dihydrofolic acid. Its effect is bacteriostatic.

Trimethoprim inhibits dihydrofolic acid reductase, which converts dihydrofolic into tetrahydrofolic acid.

The effect of trimethoprim is bacteriostatic and in combination with sulphonamides it is bactericidal.

Sulphonamides and trimethoprim thus cause a successive blockage of two enzymes that play an important role in the metabolism of bacteria and protozoa. Their effect is synergistic.

Bacterial resistance to trimethoprim and to sulphonamides can be mediated via 5 main mechanisms: (1) changes in the permeability barrier and/or efflux pumps, (2) naturally insensitive target enzymes, (3) changes in the target enzymes, (4) mutational or recombinational changes in the target enzymes, and (5) acquired resistance by drug-resistant target enzymes.

Pharmakinetics

Following oral administration, trimethoprim and sulfamethoxazole are rapidly and almost completely absorbed from the gut. The bioavailability of sulfamethoxazole is slightly higher than that of trimethoprim. It is distributed to all tissues except the brain. The highest concentrations can be found in the lungs, the liver and the kidneys.

Sulphonamides are metabolised in various ways. The degree of acetylation, hydroxylation and glucuronidation depends, among other things, on the species and age of the animal. Trimethoprim is metabolised to a large extent in the liver. Important metabolic pathways are O-methylation, N-oxidation in the ring structure and alpha hydroxylation. Sulfamethoxazole and trimethoprim are primarily excreted through the kidneys.

Indication

Treatment of bacterial diseases susceptible to sulfamethoxazole and trimethoprim

Target animals: piglets, chickens

Effective bacterial species: *E-coli*

Indication: Colibacillosis, diarrhea

Mode of Administration

Orally

Recommended Dosage

The following doses are mixed with feed and administered orally once a day for 5 to 7 days.

Usage	Volume
Feed mixing	500g of this product / administration per ton of feed

Contraindications

Do not use in animals suffering from severe liver or kidney disease, oliguria or anuria.

Do not use in animals with impaired haematopoietic systems.

Do not use in case of known hypersensitivity to sulphonamides or trimethoprim or any of the excipients.

Warnings and Precautions

Do not administer to the following animals

1. Do not use in animals that have had shock or hypersensitivity reactions to this drug and other sulfa antibiotics.
2. Do not use on pigs more than 4 months old

Interactions with Other Medicaments

Do not combine with other veterinary medicinal products.

Pregnancy and Lactation

The safety of the veterinary medicinal product during pregnancy, lactation or lay has not been established.

Adverse Effects/Undesirable Effects

A diminished water intake in chickens may occur occasionally.

Hypersensitivity reactions can occur rarely (more than 1 but less than 10 animals in 10,000 animals).

Overdose and Treatment

In chickens an acute overdose will likely not occur because the birds will be reluctant to drink the strongly concentrated drinking water (too bitter taste if above 2 litres of the veterinary medicinal product per 1000 litres drinking water). Chronic overdose in chickens will result in a strongly diminished water- and feed intake and retarded growth.

Withdrawal Period(s)

Pigs: 63 days (when slaughtered for food)

Chicken: 14 days

(* Prohibition of use of laying chickens)

Storage condition

Store in a cool dry place, protected from sunlight. Do not store above 30°C.

Keep out of reach of children/Jauhkan dari kanak-kanak.

Shelf life

Shelf life as packaged for sale: 12 months

Shelf life after reconstitution into feed: 24 hours

Pack size(s)

Plastic bag of 5kg, 10kg & 20kg

Product Registration Holder/ Manufacturer

Osmosis Nutrition Sdn. Bhd. (594876A)

Lot 16014(P.T. 24341), Jalan Nilam 3,

Bandar Nilai Utama,

71800 Nilai, Negeri Sembilan, Malaysia.

Tel: +606-799 9090 Fax: +606-799 9155

Date of revision of Package Insert

25/08/2021