

Treatment of Overdose:

In cases of overdose, methods of reducing the absorption of ingested drug are important. Gastric lavage is essential even if several hours have elapsed. Prompt administration of 50g activated charcoal and 500ml iced mannitol 20% by mouth, may reduce absorption. If the history suggests that 15g Paracetamol or more has been ingested, administer one of the following antidotes:

Acetylcysteine 20% i.v.: Administer intravenously, 20% acetylcysteine immediately without waiting for positive urine test or plasma level results: initial dose of 150mg/kg over 15 minutes, followed by continuous infusion of 50mg/kg in 500ml 5% glucose/dextrose over 4 hours and 100mg/kg in 11.5% glucose/dextrose over 16 hours;

OR

Oral Methionine: 2.5g immediately followed by three further doses of 2.5g at four hourly intervals. For a 3 year old child, 1g methionine every four hours for four doses has been used;

OR

Oral Acetylcysteine 5%: 140mg/kg as a loading dose, then 70mg/kg every 4 hours for a total of 17 maintenance doses. If more than ten hours have elapsed since the overdosage was taken, the antidote may be ineffective.

Interactions with Other Medicaments:

The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

Storage Condition:

Store in a dry place (below 30°C). Protect from light.

Pack Size:

Blister pack of 10x10's, 50x10's and 100x10's.

Product Registration Number:

MAL

Further information can be obtained from your doctor or pharmacist.

ROYCE®

Product Holder/ Manufactured by:

ROYCE PHARMA MFG SDN. BHD. 65043-X
PT 1663, Nilai Industrial Estate,
71800 Nilai, Negeri Sembilan, Malaysia.

ROYCE®

Redamol Tablet 650mg

(Paracetamol – Analgesic & Antipyretic)

Presentation:

White to off white oblong with score line on one side and plain on the other side.

Content:

Each tablet contains: Paracetamol 650 mg.

Indication:

Indicated for the relief of painful disorders such as headache; dysmenorrhea conditions involving musculoskeletal pain, myalgias and neuralgias. It is also indicated as an analgesic and antipyretic in conditions accompanied by discomfort and fever, such as the common cold and viral infections. It is an effective analgesic after dental work and tooth extraction and in teething.

Pharmacological Information:

Paracetamol is a centrally acting analgesic and antipyretic with minimal anti inflammatory properties.

Analgesic - The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (specifically cyclooxygenase (COX)-2) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation. The peripheral action may also be due to inhibition of prostaglandin synthesis or to inhibition of synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation.

Antipyretic- Paracetamol act centrally on the hypothalamic regulating center to produce peripheral vasodilatation resulting in increased blood flow through the skin, sweating and heat loss.

Paracetamol reduces fever by inhibiting the formulation and release of prostaglandins in the CNS and by inhibiting endogenous pyrogens at the hypothalamic thermoregulatory centre.

C2ff2ólyy 2WfH fRýYláuufó2yLWfHfóSfY2fH fWfHfRfE f0á2W0SfRf

tHfHfóSfY2fH f0á2Wf2y' úf1Sá WfH0S YfHyfE fy úK0S áYfHfE fyúSáúyS fYfR úK0Sf2WfS úK0S WfH0S 2f f0á2Wf2yYfE fá fRfSfYfRfYf 2y úK0S WfH0S 2f f0á2Wf2y fYfWf2yYfE Wf Kfá oSfYáK2óy' úKfHf RfWf2á f0Kf0K RfHfE f0á2Wf2y fYfWf2yYfE f0á2 RfHfE úK0S f0á2Wf2yYfE 2f WfHfHfóSfY2fE f0Kf0SfHá YfSf2f0á2Wf2YfRfS f0f RfWf2á f0Kf0K fyúSfHfSá úK0S WfH0S 2f f0á2Wf2y fYfWf2yYfE f00SfHfHfSá f0á2Wf2yYfE 2f úK0S fYfHfSáúWf úKf2Wf2K úK0S ú2WfHf fYf2Wf2y f0á2Wf2yRfS fR2áSá y2yú fyúSfHfSá

The presence of food in the stomach has also been reported to reduce the rate of absorption of paracetamol . Alterations in gastric pH have no appreciable effect on paracetamol absorption.

During absorption , the amount of paracetamol which is inactivated is negligible and it has been shown that paracetamol does not effect gastric mucosal permeability and does not produce mucosal bleeding.

tSf1f WfHámf 02y0fYfWf2yáWf2yá HfS WfSf0KfSf m K2WfHf HfSfWf f0á2Wf2yáWf2yá CkS WfHáYf KfHfWfSf fá m ú2o K2WfHf

tHfHfóSfY2fH WfSfYfHfSá úK0S WfHfYf fYfR fá WfSáSfYf fy WfSfáWf YfWf 2f K2yYfYf tHfHfóSfY2fH fá YfSfHf0á2fSf oE úK0S YfWf2á2YfHf fYfEYfS áááúSfY 2f úK0S Wf0SfWf CkHá YfSfHf0á2Yf fá YfHyfE ú2 úK0S f0Wf2Wf2YfRfS fYfR ááWfKfHfS 02yWf2YfHfSá f00á2yWfYfYf f2Wf WfWf2EYfHfSfE nqf: fYfR Hcf: 2f úK0S fy3SáúSfR fR2áS WfSáWfSfWf0SfE f0á2Wf2y ncf: fá SÉ0WfSfR fá f WfSfS WfHfHfóSfY2fE HfKfSfY YfYf2y WfHfKfHfEá fyúWf0KfS WfWf2Wf2yYfE 2f f0fSf0K2f fRfWf2Wf2Sá fYfR 0ááúSfYS 02yWf2YfHfSá WfHf f0á2Wf2yWf2ySfWf tHfHfóSfY2fE SÉ0WfSfY2fH fá WfWfRf fYfR 200Wf2á WfHf WfSfYf

52áSf3f fYfR fRýYláuufó2yLWfHf

Adults and children aged 12 years and over: One tablet every 4-6 hours (not more than 6 tablets in a day).

Contraindications:

Hypersensitivity to paracetamol or any of the other ingredient/components of the product. Severe active hepatic impairment.

Warning and Precautions:

TKfá WfSfWfHfHf2y' 02y0fHyá fWf1f9efáHf0f 52 y2yú úf1Sf fYfE 2úKfSf WfHfHfóSfY2fE 02y0fHyáYf YfSfRf0fYfSá Hf úK0S áfYfSfYfHfSf

Allergy alert:

Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

- Keep out of reach of children.

- Do not take if allergic to paracetamol.

- Patients should contact their health care provider if symptoms persist (if the pain lasts for more than 10 days, if there is redness or fever lasts more than 3 days).

- fHfSfYfá áK2WfRf 02y0fHf0f úKfSf KfSfHfKf f0HfSf WfWf2WfRfSf fy ááYfWf2Yfá WfSfWfáWf Wf úK0S WfHyf fHfáá f2Wf Y2WfSf úKfYf m n fHfEá fYf úKfSfSf fá fSfYfSáá 2Wf fSf0Sf fHfáá Y2WfSf úKfYf o RfHfá

- tHfHfóSfY2fHf áK2WfRf oS f0WfSf f0WfKf f0HfSf ú2 WfHfSfYfá f0WfKf KfSfHfWf0f fYfWfHfYfSfYf tHfHfóSfY2fHf 20SfR2áS YfHf KfHYf úK0S Wf0SfWf
f 52 y2yú SÉ0SfRfSf Wf02yYfSfYfRfSf fR2áSf f Wf áK2WfRf oS f0WfSf f0WfKf f0HfSf ú2 WfHfSfYfá f0WfKf f02K2f fRfSfYfRfSfYf0f

- tHfHfóSfHf2f WfWf2WfRfSá ááYfWf2YfHfWf WfSfSf 2yWfE fRfRfWf2YfHf úKfSfWfE ú2 WfSfHf úK0S f0HfSá 2f úK0S WfHyfWf fSf0SfWf áK2WfRf oS fyáWfWf2WfRf f0KfSfYfSf0ááHfWf

9ffSf0fá 2y f0áWfE ú2 5Wf0Sf fYfR f áS fHfKfYfáYf

Wf fá WfYf0fSfE ú2 fYfWfHf f WfHfSfYfá f0áWfE ú2 Wf0Sf 2Wf ááSf YfHfKfYfáYf

WfSfYfYfE fYfR fHf0fWf2yLWfHf

f áS fy WfSfYfYfE

f /2yáWfSfRfSf ú2 oS úK0S fYfHfSáúWf 2f 0K2f0S fy WfSfYfYfYf WfHfSfYfá f f WfK2WfKf Wf 0f2ááSá WfHfSfYfE WfHfHfóSfY2fHf á 02yáWfSfRfSf ú2 oS áfSf fyáWfYfHf úKfSfWfSfWf0f fR2áSá f2Wf áK2WfWfSfYf WfáS fá f YfYf2y fYfHfSáWf0KfYfWfWfWfSfWf0f fy WfSfYfYfYf0f

f áS fy fHf0fWf2yLWfHf

f 9É0WfSfRfSf fy WfSfáWf YfWf

f á HfSfYfHf fy3Sáú2y 2f WfHfHfóSfY2fHf fyáWfYfHf úKfSfWfSfWf0f fR2áSá fR2áSá y2yú WfWfSfHf ú2 WfSáSfYf0f f WfáLf ú2 úK0S yWfáWfYf fyfYf0f

fRfSf 9ffSf0fáYf

fRfSfWfSf fSfSf0fá 2f WfHfHfóSfY2fHf HfSf WfHfSf fYfR WfáWfHfE YfWfRf HfK2Wf2Kf KfYfSf Hf2E2f0fHf WfSf0fWf2yá Kf0S oSfSfYfWf2Wf2SfRf /WfWfYfSf2Wfá KáUfSáYfáWfWfWfE WfSf0fWf2yá fyúWfRfYfá áfYf WfáKfSá fYá2fRfSfYfHf f0Sf0fYá Wf2Kfá2y fEYfRfYfSfKf C2E0f 9WfRfSfYfHf bSfWf2fEáá fKf0S oSfSfYfWf2Wf2SfRf

Symptoms and Treatment of Overdose:

Symptoms:

Toxic symptoms include vomiting, abdominal pain, hypotension and sweating. The most serious adverse effect of acute overdose of paracetamol is a dose-dependent, potentially fatal hepatic necrosis. Clinical and laboratory evidence of hepatotoxicity may be delayed for up to one week. Major manifestations of liver failure such as jaundice, hypoglycemia and metabolic acidosis may take at least 3 days to develop.