

Watsons OTC_Pain & Fever Relief EXTRA Tablet insert (MY version)

23-02-2022(R9)

98 mm

140 mm

Watsons Pain & Fever Relief EXTRA Tablet

Name and Strength of Active Substance(s)
Paracetamol 500mg
Caffeine 65mg

Product Description

Watsons Pain and Fever Relief EXTRA Tablet is a white, modified capsule-shaped, 0.662" x 0.300" in diameter, convex, bisected line on one side and plain on the other side.

Pharmacodynamics

Paracetamol
Paracetamol is a centrally acting analgesic and antipyretic with minimal anti-inflammatory properties

Analgesic

The mechanism of analgesic action has not been fully determined. Paracetamol may act predominantly by inhibiting prostaglandin synthesis in the central nervous system (CNS) (specifically cyclooxygenase (COX)-2) and, to a lesser extent, through a peripheral action by blocking pain-impulse generation.

The peripheral action may also be due to inhibition of prostaglandin synthesis or inhibition of the synthesis or actions of other substances that sensitize pain receptors to mechanical or chemical stimulation.

Antipyretic

Paracetamol acts centrally on the hypothalamic heat-regulating centre to produce peripheral vasodilatation resulting in increased blood flow through the skin, sweating and heat loss.

Paracetamol reduces fever by inhibiting the formulation and release of prostaglandins in the CNS and by inhibiting endogenous pyrogens at the hypothalamic thermoregulatory centre.

Caffeine

Caffeine is a CNS stimulant. It stimulates all levels of the CNS, although its cortical effects are milder and of shorter duration than those of amphetamines.

Analgesia adjunct

Caffeine constricts cerebral vasculature with an accompanying decrease in the cerebral blood flow and in the oxygen tension of the brain. It is believed that caffeine helps to relieve headache by providing more rapid onset of action and/or enhancing pain relief with lower doses of analgesic.

The combination of paracetamol and caffeine is a well-established analgesic combination.

Pharmacokinetics

Paracetamol is rapidly and almost completely absorbed from the gastro-intestinal tract. It is relatively uniformly distributed throughout most body fluids and it exhibits variable protein binding. Excretion is almost exclusively renal in the form of conjugated metabolites.

Caffeine is absorbed readily after oral administration, maximal plasma concentrations are achieved within one hour and the plasma half-life is about 3.5 hours. 65-80% of administered caffeine is excreted in the urine as 1-methyluric acid and 1-methylxanthine.

Indications

Paracetamol-caffeine is indicated for the treatment of most painful and febrile conditions eg, headache, including migraine, toothache, rheumatic pain and menstrual pain. Relief of the symptoms of colds, influenza and sore throat.

Recommended Dosage

Adults and children aged 12 years and over

Take 1-2 tablets four times daily as required. The dose should not be repeated more frequently than every 4 hours. Do not exceed 8 tablets in 24 hours.

Children under 12 years

Not recommended for children under 12 years.

Mode of Administration

Oral

Contraindications

Hypersensitivity to paracetamol, caffeine or any of the other constituents.

Warnings and Precautions

- Avoid other caffeine containing products. Too much caffeine may cause rapid heart rate, nervousness or sleeplessness.
- Ask a doctor or pharmacist before use if you have high blood pressure, glaucoma, or overactive bladder syndrome.
- DO NOT** exceed 8 tablets in 24 hours.
- DO NOT** take more than the recommended dose unless advised by your doctor. Use the smallest effective dose. Taking more than the maximum daily dose may cause severe or possibly fatal liver damage.
- DO NOT** use with other drugs containing **paracetamol**.
- Not recommended for children under 12 years.
- Care is advised in the administration of paracetamol to patients with renal or hepatic impairment. The hazard of overdose is greater in those with non-cirrhotic alcohol liver disease.
- Excessive intake of caffeine (e.g. coffee, tea and some canned drinks) should be avoided while taking this product.
- Patients should be advised to consult their doctor if their headaches become persistent.
- Keep medicine out of reach of children.

Allergy alert:

Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Interactions with Other Medications

The speed of absorption of paracetamol may be increased by metoprolamide or domperidone and reduced by colestyramine. The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

KNIFE GUIDE INSERT

Dimension : 98 x 140 mm

Pregnancy and Lactation

Paracetamol-caffeine is not recommended for use during pregnancy due to the possible increased risk of lower birth weight and spontaneous abortion associated with caffeine consumption.

Caffeine in breast milk may potentially have a stimulating effect on breast fed infants.

Due to the caffeine content of this product, it should not be used if you are pregnant or breast feeding.

Adverse Effects / Undesirable Effects

Adverse reactions of paracetamol-caffeine are rare when taken in recommended doses, but some undesirable effects such as skin rash or itching have been reported, sometimes with breathing problems or swelling of the lips, tongue, throat or face. Bruising or bleeding.

Adverse Effect of Paracetamol

The side effects are rare and have been reported according to the body system; Thrombocytopenia and agranulocytosis (blood and lymphatic system disorders), anaphylaxis, cutaneous hypersensitivity reactions including skin rashes, angioedema and Stevens Johnson Syndrome/toxic epidermal necrolysis (immune system disorders), bronchospasm (respiratory, thoracic and mediastinal disorders), hepatic dysfunction (hepatic effects).

Adverse Effect of Caffeine

Nervousness and dizziness have been reported for the CNS.

When the recommended paracetamol-caffeine dosing regimen is combined with dietary caffeine intake, the resulting higher dose of caffeine may increase the potential for caffeine-related adverse effects such as insomnia, restlessness, anxiety, irritability, headaches, gastrointestinal disturbances and palpitations.

Overdose and Treatment

Paracetamol

Symptoms

Abdominal pain. Liver damage may become apparent 12 to 48 hours after ingestion. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning, hepatic failure may progress to encephalopathy, haemorrhage, hypoglycaemia, cerebral oedema and death. Acute renal failure with acute tubular necrosis, strongly suggested by loin pain, haematuria and proteinuria, may develop even in the absence of severe liver damage. Cardiac arrhythmias and pancreatitis have been reported.

Treatment

Immediate treatment is essential in the management of paracetamol overdose. Despite a lack of significant early symptoms, patients should be referred to hospital urgently for immediate medical attention. Symptoms may be limited to nausea or vomiting and may not reflect the severity of the overdose or the risk of organ damage.

Treatment with activated charcoal should be considered if the overdose has been taken within 1 hour. Plasma paracetamol concentration should be measured at 4 hours or later after ingestion (earlier concentrations are unreliable). Treatment with N-acetylcysteine may be used up to 24 hours after ingestion of paracetamol, however, the maximum protective effect is obtained up to 8 hours post-ingestion. The effectiveness of the antidote declines sharply after this time. If required, the patient should be given intravenous N-acetylcysteine, in line with the established dosage schedule. If vomiting is not a problem, oral methionine may be a suitable alternative for remote areas outside hospital. Management of patients who present with serious hepatic dysfunction beyond 24 hours from ingestion should be discussed with relevant health care professionals.

Caffeine

Symptoms

Overdose of caffeine may result in epigastric pain, vomiting, diuresis, tachycardia or cardiac arrhythmia, and CNS stimulation (insomnia, restlessness, excitement, agitation, jitteriness, tremors and convulsions).

It must be noted that for clinically significant symptoms of caffeine overdose to occur with this product, the amount ingested would be associated with serious paracetamol-related toxicity.

Treatment

Patients should receive general supportive care (e.g. hydration and maintenance of vital signs). The administration of activated charcoal may be beneficial when performed within one hour of the overdose, but can be considered for up to four hours after the overdose. The CNS effects of overdose may be treated with intravenous sedatives.

Storage Conditions

Store in dry place at room temperature below 30°C. Do not use if foil pack is broken.

Dosage Forms and Packaging Available

In box of 20's (2 blisters x 10 tablets)

Name and Address of Manufacturer

PT. Medifarma Laboratories
Jl. Raya Jakarta - Bogor Km.33, Kelurahan Curug, Kecamatan Cimanggis, Kota Depok, Jawa Barat, Indonesia

Name and Address of Product Registration Holder

Unam Pharmaceutical (M) Sdn Bhd
Suite 803, Block F, Phileo Damansara 1, No. 9, Jalan 16/11, 46350 Petaling Jaya, Selangor Malaysia

Distributed by Watsons: Diedarkan oleh Watsons:

Watson's Personal Care Stores Sdn Bhd (289892-V)
19th Floor Wisma Chuang, 34 Jalan Sultan Ismail, 50250 Kuala Lumpur, Malaysia
Customer Service Hotline: 1 300 880 847

Date of Revision of Package Insert

01/12/2021

Black

FRONT SIDE

BACK SIDE