

**NAME AND STRENGTH OF ACTIVE SUBSTANCE:**

Acyclovir 5% w/w.

**PRODUCT DESCRIPTION:**

A smooth odourless white cream.

**PHARMACODYNAMICS:**

Acyclovir is an antiviral agent which is highly active in vitro against herpes simplex virus (HSV) types I and II and varicella zoster virus. Toxicity to mammalian host cells is low.

Acyclovir is phosphorylated after entry into herpes infected cells to the active compound acyclovir triphosphate. The first step in this process is dependent on the presence of the HSV-coded thymidine kinase. Acyclovir triphosphate acts as an inhibitor of and substrate for, the herpes-specified DNA polymerase, preventing further viral DNA synthesis without affecting normal cellular processes.

**PHARMACOKINETICS:**

Only minimal systemic absorption of acyclovir following repeated topical administration of Vivere Cream 5% w/w.

**INDICATION:**

Acyclovir cream is indicated for the treatment of herpes simplex virus infections of the skin including initial and recurrent genital herpes and herpes labialis.

**RECOMMENDED DOSAGE:**

*Adults and children:*

Vivere Cream 5% w/w should be applied five times daily at approximately four hourly intervals omitting the night time application.

Vivere Cream 5% w/w should be applied to the lesions or impending lesions as soon as possible, preferably during the earliest stages (prodrome or erythema). Treatment can also be started during the later (papule or blister) stages.

Treatment should be continued for at least 4 days for herpes labialis and for 5 days for genital herpes. If healing has not occurred treatment may be continued for up to 10 days.

**ROUTE OF ADMINISTRATION:**

Topical

**CONTRAINDICATION(S):**

Contraindicated in patients known to be hypersensitive to acyclovir, valacyclovir or any of the excipients of Vivere Cream 5% w/w.

**WARNINGS AND PRECAUTIONS:**

For External Use Only.

Vivere Cream 5% w/w is not recommended for application to mucous membranes, such as in the mouth, eye or vagina, as it may be irritant. Particular care should be taken to avoid accidental introduction into the eye.

In severely immune-compromised patients (e.g. AIDS patients or bone marrow transplant recipients) oral acyclovir dosing should be considered. Such patients should be encouraged to consult a physician concerning the treatment of any infection.

**INTERACTIONS WITH**
**OTHER MEDICAMENTS:**

No clinically significant interactions have been identified.

**TEXT SIZE IN THIS ARTWORK**

Body text size: 13.5 pt

Smallest text size: 12.5 pt

Microtext: No

**STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:****Pregnancy:**

The use of Vivere Cream 5% w/w should be considered only when the potential benefits outweigh the possibility of unknown risks however the systemic exposure to acyclovir from topical application of Vivere Cream 5% w/w is very low.

**Lactation:**

The dosage received by a nursing infant following maternal use of Vivere Cream 5% w/w would be insignificant.

**ADVERSE EFFECTS/****UNDESIRABLE EFFECTS:*****Skin and subcutaneous tissue disorders*****Uncommon**

- Transient burning or stinging following application of Vivere Cream 5% w/w
- Mild drying or flaking of the skin
- Itching

**Rare**

- Erythema
- Contact dermatitis following application. Where sensitivity tests have been conducted, the reactive substances have most often been shown to be components of the cream rather than acyclovir.

***Immune system disorders*****Very rare**

- Immediate hypersensitivity reactions including angioedema.

**OVERDOSE AND TREATMENT:**

No untoward effects would be expected if the entire contents of a 5 gram tube of Vivere Cream 5% w/w containing 250 mg of acyclovir were ingested orally. Acyclovir is dialysable by haemodialysis.

**INSTRUCTIONS FOR USE:**

For topical use only. Vivere Cream 5% w/w should be applied five times daily at approximately four hourly intervals omitting the night time application.

**STORAGE CONDITIONS:**

Store below 30°C. Do not refrigerate.

**SHELF LIFE:**

3 years.

**PACKAGING AVAILABLE:**

Each box contains 1 aluminium tube of 5 g cream.

**Manufacturer:****SIAM BHEASACH CO., LTD.**

123 Soi Chokechairuammitr, Vibhavadi-Rangsit Road, Chomphon, Chatuchak, Bangkok 10900 and 9 Soi Chokechairuammitr 3, Vibhavadi-Rangsit Road, Dindang, Dindang, Bangkok 10400, Thailand.

**Product Registration Holder,****Importer:****SPG PHARMA (MALAYSIA) SDN BHD**

Suite 1701, Block A, Damansara Intan e-Business Park, No. 1, Jalan SS20/27, 47400 Petaling Jaya, Selangor Darul Ehsan, Malaysia.

**DATE OF REVISION OF PI:**

30/11/2021

VIV-CREAM-5-04-L-ENG-MY-A0001

Copyright © 2021 Siam Bheasach. All rights reserved.

**TEXT SIZE IN THIS ARTWORK**

Body text size: 13.5 pt

Smallest text size: 12.5 pt

Microtext: No