

AMOX FORTE

Amoxicillin trihydrate 500 mg/g, water soluble powder

AMOX FORTE is an oral water soluble powder containing Amoxicillin trihydrate.

Pharmacodynamics

AMOX FORTE is an oral powder containing Amoxicillin trihydrate. Amoxicillin is a time-dependent bactericidal antibiotic which acts by inhibiting the synthesis of bacterial cell walls during bacterial replication. It inhibits the formation of bridges between the chains of linear polymers constituting the peptidoglycan cell wall of Gram positive bacteria. Amoxicillin is a broad-spectrum penicillin. It is also active against a limited range of Gram negative bacteria on which the outer layer of the bacterial cell wall is composed of lipopolysaccharide and proteins.

There are three main mechanisms of resistance to beta-lactams: beta-lactamase production, altered expression and/or modification of penicillin binding proteins (PBP), and decreased penetration of the outer membrane. One of the most important is the inactivation of penicillin by beta-lactamase enzymes produced by certain bacteria. These enzymes are capable of cleaving the beta-lactam ring of penicillins, making them inactive. The beta-lactamase could be encoded in chromosomal or plasmidic genes.

Cross-resistance is observed between amoxicillin and other penicillins, particularly with aminopenicillins. The use of extended spectrum beta-lactam drugs (e.g. aminopenicillins) might lead to the selection of multi-resistant bacterial phenotypes (e.g. those producing extended spectrum beta-lactamases (ESBLs)).

Pharmacokinetics

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids. Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

Environmental Properties

Not applicable. AMOX FORTE is not used directly to environment

COMPOSITION

Each g contains:
Amoxicillin trihydrate 500 mg
Excipients up to 1 g

Preservative : Sodium benzoate 1 mg/g

INDICATION

- Chickens, turkeys and ducks : Treatment of infections caused by bacteria susceptible to amoxicillin
- Pigs : For the treatment of pasteurellosis caused by *Pasteurella multocida* susceptible to amoxicillin

RECOMENDED DOSAGE

Species	Recommended dose
Chickens	The recommended dosage is 15 mg amoxicillin trihydrate per kg body weight per day (corresponding to 30 mg product/kg body weight/day). The total period of treatment should be for 3 days or in severe cases for 5 days
Ducks	The recommended dosage is 20 mg amoxicillin trihydrate per kg body weight per day (corresponding to 40 mg product/kg body weight/day) for 3 consecutive days
Turkey	The recommended dosage is 15-20 mg amoxicillin trihydrate per kg body weight per day (corresponding to 30 - 40 mg product/kg body weight/day) for 3 days or in severe cases for 5 days
Pigs	For the medication of pigs, the product may be administered via the drinking water. It may not be used in dry feeds. Administer in the drinking water to give 20 mg amoxicillin trihydrate per kg body weight (corresponding to 40 mg product/kg body weight/day) daily for up to 5 days. Prepare the solution by carefully mixing the product in the requisite quantity of fresh potable water immediately before use. The dose should be administered at approximately 24 hourly intervals for up to 5 days. Any medicated water which is not consumed within 24 hours should be discarded and the medicated drinking water replenished. In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being on treatment

Instruction for use

For administration in drinking water.
Prepare the solution with fresh potable water immediately before use.
Any medicated water which is not consumed within 24 hours should be discarded and the medicated drinking water replenished.
In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated.

Use the following formula in order to calculate the concentration of the product (mg) per litre of drinking water:
$$\frac{\text{Dose (mg product per kg body weight per day)} \times \text{mean body weight (kg) of animals to be treated}}{\text{mean daily water consumption (litre) per animal per day}} = \text{___ mg product per litre drinking water}$$

To ensure a correct dosage, body weight should be determined as accurately as possible to avoid under dosing. The uptake of medicated water depends on the clinical condition of the animal. In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted taking into account water intake.

The maximum solubility of the product was only demonstrated at 5 g/L at 20°C. Below 20°C and above 5 g/L, the product cannot be satisfactorily dissolved. For stock solutions and when using a proportioner, take care not to exceed the maximum solubility which can be achieved under the given conditions. Adjust the flow rate settings of the dosing pump according to concentration of the stock solution and water intake of the animals to be treated.
After the end of the medication period, the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance.

CONTRAINDICATION

- Do not administer to rabbits, guinea pigs, hamsters, horses, gerbils or any other small herbivore
- Do not use in animals with known hypersensitivity to penicillins or other β-lactam antibiotics or to the excipient
- Do not administer to animals with renal disease including anuria or oliguria

WARNING AND PRECAUTION

Pigs: The uptake of medication by animals may be altered as a consequence of illness. In case of insufficient water uptake, animals should be treated parenterally instead using a suitable injectable product prescribed by the veterinarian.

Special precautions for use in animals

- Not effective against beta-lactamase producing organisms
- Official, national and regional antimicrobial policies should be taken into account when the product is used
- Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria
- Use of the product deviating from the instructions given may increase the prevalence of bacteria resistant to the amoxicillin and may decrease the effectiveness of the treatment

Special precautions to be taken by the person administering the veterinary medicinal product to animals

- Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious
- Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations
- Handle this product with great care to avoid exposure, taking all recommended precautions
- If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention
- Avoid inhalation of dust. Wear either a disposable half-mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143
- Wear gloves during preparation and administration of medicated water or liquid feed
- Wash any exposed skin after handling the product or medicated water or feed. Wash hands after use

INTERACTION

The product should not be administered with antibiotics that have a bacteriostatic mode of action, such as tetracyclines, macrolides, sulphonamides.

PREGNANCY AND LACTATION

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.
The safety of the veterinary medicinal product has not been established during pregnancy or lactation in sows. Use only accordingly to the benefit-risk assessment by the responsible veterinarian

ADVERSE EFFECT

Penicillins and cephalosporins may cause hypersensitivity reactions which may occasionally be serious. Rarely, gastrointestinal tract signs associated with alteration of the intestinal flora (for example, loose stools, diarrhea) may occur

OVERDOSE AND TREATMENT

No problems with over dosage have been reported. Treatment should be symptomatic and no specific antidote is available

WITHDRAWAL PERIOD

Meat and offal:

- Chickens : 1 day
- Ducks : 9 days
- Turkeys : 5 days
- Pigs : 2 days

STORAGE

Store at temperature below 30°C in a tightly closed container, in a dry and cool place, protected from direct sunlight

PACKING

Sachet : 250 g

ANTIBACTERIAL

For animal use only

Controlled medicine/ Ubat terkawal
Keep out of reach of children
Jauhkan daripada capaian kanak-kanak

To avoid microbial contamination, unused portions of the product must be discarded within 24 hours after first opening of the container or after dilution

Disposal of containers- in accordance to federal, state and local environment control regulations

Manufactured by : PT Medion Farma Jaya, Jl Raya Batuajar No 29, Cimagreme, Ngamprah, Bandung Barat – Indonesia

Product registration holder : Sum Soon Agricultural Co. Sdn. Bhd., Lot 42, Bandaran Berjaya, P.O. BOX 12061, 88822 Kota Kinabalu, Sabah, Malaysia

Date of revision : March 2023

MAL number :

