

ERYTROSURE 50% POWDER

Description and Composition

ERYTROSURE 50% POWDER is a white to off-white powder. Each gram contains 500mg Erythromycin (as thiocyanate) as its active ingredient.

Pharmacodynamics

Erythromycin is a macrolide antibiotic with a bacteriostatic action against a wide range of bacteria, although it also has a bactericidal action against certain pathogenic organisms. It is highly active against gram-positive bacteria and at low concentrations it inhibits many other types of microorganisms. In addition to the gram-positive bacteria, most strains of *Neisseria* and *Haemophilus* are sensitive as are some strains of *Bordetella*, *Brucella*, *Pasteurella*, *Listeria*, *Actinomyces*, *Mycoplasma*, *Rickettsia*, certain large viruses and *Treponema pallidum*.

Erythromycin exerts its effect only against multiplying bacteria. It inhibits protein synthesis by binding to 50 S ribosomal subunits of sensitive microorganisms. Certain resistant micro-organisms with mutational changes in components of this subunit of the ribosome fail to bind the drug.

Pharmacokinetics

Erythromycin is absorbed rapidly when administered orally and diffusion occurs into most tissues and fluids. Elimination of erythromycin occurs primarily through hepatic metabolism and the remainder is excreted in active form in the urine and bile.

Indication

For the treatment of Chronic Respiratory Disease caused by *Mycoplasma* infection.

Recommended Dose

For oral administration via the drinking water.

The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of erythromycin has to be adjusted accordingly.

Mix 33g Erytosure 50% Powder per 64 litres of drinking water and administer continuously for 1 to 5 days, depending on response (approximately 25.5mg Erythromycin per kg body weight per day).

Slowly add 33g to not less than 1.07 litres of clean, cold drinking water, stirring continuously until completely dissolved. This solution should then be added to more drinking water to make up a total volume of 64 litres.

Route of Administration

For oral administration via drinking water.

Contraindications

Any medicated drinking water which is not consumed within 24 hours should be discarded.

Warning and Precautions

Special warnings for each target species:

None.

Special precautions for use

Special precautions for use in animals:

The water system to which the product is added must be in good working order and must be free of dust, algae or other particulate matter.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

Due to a lack of information on the fate and effects of erythromycin in the environment veterinary surgeons prescribing this product under the cascade should satisfy themselves that there will be no adverse impact on the environment.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Do not eat, drink or smoke while preparing the diluted formulation or while animals are being treated. In case of contact with eyes, rinse immediately with clean water and seek medical advice if irritation persists. Wash hands and exposed skin after use.

Interaction with Other Medicaments

None known.

Pregnancy and Lactation

Not applicable.

Side Effects

None known.

Symptoms and Treatment of Overdose

None known.

Storage Condition

Store below 30°C.

Shelf life

3 years.

To be used within 24 hours after dilution in drinking water.

Withdrawal Periods

Chickens: 6 days

Eggs: 6 days

Packing

100g, 200g, 500g, 1kg.

Revised date: 20.6.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com