

MOXILA 15% INJECTION

Description and Composition

MOXILA 15% INJECTION is an off white to grayish yellow oily suspension. Each ml contains 150mg of Amoxicillin (as trihydrate) as its active ingredient and 1mg Chlorocresol as its preservative.

Pharmacodynamics

Amoxicillin predominately inhibits cell wall synthesis in susceptible bacteria. Amoxicillin has a unique mode of action which directly and irreversibly disrupts existing cell wall peptidoglycan rather than newly forming peptidoglycan of the divisory septal wall as with other members of the penicillin family.

Pharmacokinetics

Amoxicillin is primarily distributed to the extracellular compartment. Its distribution in the tissues is facilitated by the fact that it binds little to plasma proteins. Concentrations in lung, pleural and bronchial tissues are identical to those found in plasma. Amoxicillin diffuses into pleural and synovial fluid as well as into lymphoid tissues.

A small proportion of amoxicillin (approximately 20%) is biotransformed in the liver by hydrolysis of the β -lactam ring, producing inactive penicilloic acid.

Amoxicillin is mainly excreted in active form by the kidneys, then through the bile and through milk.

Indication

In vitro Amoxicillin is effective against a wide range of Gram-positive and Gram-negative bacteria which include: *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus* species, *Salmonella* species, Staphylococci and Streptococci.

Not effective against beta-lactamase producing organisms.

Suitable for the control of infections due to susceptible microorganisms in cattle, sheep, pigs, dogs and cats where a single injection giving prolonged activity is required. It may also protect from secondary bacterial invasion due to sensitive organisms in cases where bacteria are not the initial cause of the disease.

Indications include infections of: Alimentary tract, respiratory tract, skin and soft tissue, urogenital tract and in prevention of post-operative infection (treat before surgery).

Recommended Dose

Cattle, sheep and pigs - By intramuscular injection only.

Dogs and cats - By subcutaneous or intramuscular injection.

The recommended dosage rate is 15mg per kg body weight (equivalent to 1ml per 10kg body weight), repeatable if necessary after 48 hours. Massage the injection site.

If dose volume exceeds 15ml in cattle and 4ml in sheep and pigs, it should be divided and injected into two or more sites. Shake well before use.

Animal	Weight (kg)	Dose volume (ml)
Cattle	450kg	45.0ml
Sheep	65kg	6.5ml
Pigs	150kg	15.0ml
Dogs	20kg	2.0ml
Cats	5kg	0.5ml

A separate injection site should be used for each administration. Use a dry sterile needle and syringe. An appropriately graduated syringe must be used to allow accurate administration of the required dose volume. This is particularly important when injecting small volumes. To ensure the correct dosage, body weight should be determined as accurately as possible to avoid underdosing or overdosing. Swab the

septum before removing each dose. If no distinct clinical response is seen after the second treatment, a check of the diagnosis and eventually a change of treatment are required.

Route of Administration

Cattle, sheep and pigs - By intramuscular injection only.

Dogs and cats - By subcutaneous or intramuscular injection.

Contraindication

This product is not suitable for intravenous or intrathecal use.

This product should not be administered to rabbits, hamsters, gerbils or guinea pigs.

Not for use in known cases of hypersensitivity to penicillins or cephalosporins.

Warning and Precautions

Special warnings for each target species:

No special warnings.

Special precautions for use:

Special precautions for use in animals:

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal.

If this is not possible, therapy should be based on local epidemiological information.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Care should be taken to avoid accidental self-injection. In the case of accidental self-injection, seek medical advice immediately.

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross-reactions to cephalosporins and vice versa. Allergic reactions to these substances may occasionally be serious.

1. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations.

2. Handle this product with great care to avoid exposure, taking all recommended precautions.

3. If you develop symptoms following exposure such as skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing, are more serious symptoms and require urgent medical attention.

Wash hands after use.

Interaction with Other Medicaments

It is not generally recommended to use bactericidal and bacteriostatic antibiotics at the same time.

Pregnancy and Lactation

Can be safely administered during pregnancy and lactation.

Side Effects

In rare cases, hypersensitivity reactions such as urticaria anaphylaxis shock can occur after use. In the case of allergic reactions, treatment should be discontinued and a symptomatic treatment should be initiated. In very rare cases, local tissue reactions such as swelling and pruritus may result from the use of amoxicillin.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s))
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports)

Symptoms and Treatment of Overdose

Penicillins have a wide safety margin.

Storage Condition

Store below 30°C.

Shelf life

3 years. Following withdrawal of the first dose, use the product within 24 hours. Any unused material should be discarded.

Withdrawal Periods

Cattle: Meat and offal: 39 days

Milk: 108 hours (4.5 days)

Pigs: Meat and offal: 42 days

Sheep: Meat and offal: 29 days

Milk: Not authorised for use in sheep producing milk for human consumption.

Incompatibilities

None known

Packing

20ml, 50ml, 100ml, 250ml.

Revised date: 13.4.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com