

PASTORAL STAR CTC 15% w/w Granule

Name and Strength of Active Substance: 150 mg of Chlortetracycline Hydrochloride per g product

Product Description: Brown to dark brown granule, not lumpy or moldy, no unpleasant odor.

Pharmacodynamics/ Pharmacokinetics/ Environmental Properties:

Chlortetracycline hydrochloride is a predominantly bacteriostatic antibiotic, interfering with bacterial protein synthesis of the rapidly growing and reproducing bacterial cell. Chlortetracycline has a broad spectrum of activity, including Gram-positive aerobes, Gram-negative anaerobes and Mycoplasmas. Resistance is known to occur in respiratory pathogens of pigs and cross-resistance occurs between chlortetracycline and other tetracyclines. The Clinical and Laboratories Standards Institute (CLSI) breakpoints established for tetracyclines are as follows: Organisms other than streptococci: S: $\leq 4\mu\text{g/ml}$, I: $8\mu\text{g/ml}$; R: $\geq 16\mu\text{g/ml}$.

Swine oral administration, maximum blood levels are achieved within approximately 2 - 8 hours. Steady state plasma concentrations of chlortetracycline are maintained throughout the twice daily seven day treatment period.

Indication and Recommended Dosage:

Chicken:

- 1) Control of infectious synovitis caused by Mycoplasma synoviae susceptible to chlortetracycline. 110-220 grams per tonne of feed. Feed continuously for 7 to 14 days.
- 2) Reduction of mortality due to Escherichia coli infections susceptible to chlortetracycline. 550 grams per tonne of feed. Feed for 5 days.
- 3) Control of chronic respiratory disease and air sac infection caused by Mycoplasma gallisepticum and Escherichia coli susceptible to chlortetracycline. 220-440 grams per tonne of feed. Feed continuously for 7 to 14 days.

Swine:

- 1) Treatment of bacterial enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia caused by Pasteurella multocida susceptible to chlortetracycline. Feed approximately 440 grams per tonne of feed, varying with body weight and feed consumption to provide 22 milligrams per kg of body weight per day in feed. Feed for not more than 14 days.
- 2) Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group E Streptococci susceptible to chlortetracycline. 55-110 grams per tonne of feed.

Swine, breeding: Control of leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by Leptospira pomona susceptible to chlortetracycline. 440 grams per tonne of feed. Feed continuously for not more than 14 days.

Mode of Administration: Orally into feed

Contraindications: Do not use in animals with known hypersensitivity to tetracycline. Do not use in animals with severe liver and renal disorders.

Warnings & Precautions:

Special precautions for use in animals: The product is efficient only against bacterial strains most sensitive to chlortetracycline. Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about the susceptibility of the target bacteria. Inappropriate use of the product may increase the prevalence of bacteria resistant to chlortetracycline and may decrease the effectiveness of treatment with related substances, due to the potential for cross-resistance. Long term use of this product is not recommended as it may lead to the development of bacterial resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals: Handle this product with care to avoid exposure when adding to feed and administering medicated feed to animals. Take adequate measures to avoid dust formation when adding the product to feed. Those handling the product should do so in a mechanically ventilated area. Wear either a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143. Direct contact of the product with the skin, eyes and mucous membranes should be avoided. Wear protective gloves, overalls and approved safety glasses. In case of accidental exposure, wash area immediately with water. Do not smoke, eat or drink when handling the product. Hands and exposed skin should be washed thoroughly after use.

Interactions with Other Medicaments: This product is not recommended for concurrent administration with any other oral medication. Do not incorporate the product in feed overloaded with polyvalent cations such as Ca^{2+} and Fe^{3+} because the formation of chlortetracycline complexes with these cations is possible. Do not administer together with antacids, kaolin and iron preparations and in conjunction with bactericidal antibiotics like beta-lactams. The product should not be used in case of known resistance to other tetracyclines.

Statement on usage during pregnancy and lactation: The use is not recommended during pregnancy or lactation. The treatment of pregnant animals with chlortetracycline may result in adverse effects on skeletal and tooth development in the foetus. Therefore, the product should be used only in pregnant sows according to the benefit/risk assessment of the responsible veterinarian.

Adverse Effects/Undesirable Effects: Chlortetracycline toxicity is low. If digestive disturbances do occur, treatment should be discontinued. On rare occasions (more than 1 but less than 10 animals in 10,000 animals) the following adverse reactions may occur: allergic reactions and photosensitivity; gastrointestinal disorders; disorders of the liver and the kidneys. If suspected adverse reactions occur, treatment should be discontinued.

Overdose and Treatment: Do not exceed the stated dose. Chlortetracycline toxicity is low. If digestive disturbances do occur, treatment should be discontinued.

Withdrawal Period: Chicken: 24 hours before slaughter when fed at 550 grams per tonne.

Swine: zero days before slaughter.

Storage Condition: Sealed and stored in a cool (<25°C) and dry place without sunlight.

Dosage Form: Granule

Packaging available: 25kg/bag

Name and Address of Manufacturer: Jinhe Biotechnology Co., Ltd. No. 71 Xiping Road, Tuoketuo County, Hohhot 010200, Inner Mongolia, P.R. China

Name and Address of Marketing Authorisation Holder: Depco United Sdn Bhd, 2433 Mk 1, Tingkat Perusahaan 6, Prai Industrial Estate, 13600 Prai, Penang, Malaysia.

Date of Revision of Package Insert: 17 October 2023