

COSULTRIM PLUS 75% W/W WATER SOLUBLE POWDER

STRENGTH OF ACTIVE SUBSTANCE COMPOSITION:

Per 100gm

Trimethoprim..... 12.50gm

Sulfachloropyridazine Sodium62.50 gm

DESCRIPTION:

A buff or light yellow color powder.

DOSAGE FORM:

Powder to be dissolved in water

PHARMACODYNAMICS:

The antibacterial activity of the veterinary medicinal product is based on a double sequential inhibition of the microbial synthesis of tetrahydrofolic acid. Trimethoprim is an antibacterial diaminopyrimidine derivative having bactericidal and synergistic activity with sulfonamides. Sulfachloropyridazine is a sulfa antibacterial drug that works by preventing bacterial growth. The combination of the two active substances has a synergistic effect, which influences the dosage (reduction in doses of active substances) and the bactericidal effect in a positive way and broadens the spectrum of activity. Furthermore, the appearance of bacterial resistance is slowed.

In chickens the veterinary medicinal product, administered at a dose of 200 mg/kg body weight, is active against respiratory and septic infections by wild-type *E. coli* and against septic infections by *P. multocida* and *S. aureus*.

Resistance develops slowly through transferable elements and results from chromosomal mutation as a result of one or more of the mechanisms described below:

- permeability barrier and/or active efflux;
- insensitive natural target enzymes;
- adjustable modifications of the target enzyme;
- mutational or recombinational modifications of the target enzyme;
- resistance acquired by target enzymes resistant to the veterinary medicinal product

PHARMACOKINETICS:

The pharmacokinetic parameters in chickens after a single administration of the veterinary medicinal product at the recommended daily dose are summarized in the table below, as well as the steady state plasma concentrations (C_{ss}) by continuous medication in drinking water. Significantly higher plasma concentrations are achieved with sulfachloropyridazine.

	Triméthoprim	Sulfachloropyridazine
C _{max} (µg/ml)	0,52	18,65
AUC _{24h} (µg.h/ml)	2,14	103,4
T _{max} (h)	1,55	1,32
T _{1/2el} (h)	1,41	2,60
C _{ss} (µg/ml)	0,089	4,31

Protein binding in plasma reaches 24.1% for trimethoprim and 35.1% for sulfachloropyridazine.

These results translate into an effective C_{ss} (steady state) of 0.068 µg/ml for trimethoprim and 2.797 µg/ml for sulfachloropyridazine. Both active components are absorbed approximately equally quickly and excreted relatively quickly

TARGET SPECIES:

Poultry (Chicken)

INDICATIONS:

Treatment of infections caused by germs sensitive to trimethoprim/sulfachloropyridazine, taking into account the antibiotic's ability to reach the site of infection in effective concentrations

RECOMMENDED DOSAGE :

32 mg COSULTRIM PLUS 75% W/W WATER SOLUBLE POWDER (equivalent with 24 mg active substance) per kg body weight per day.

Mode of administration :

Chicken: mix COSULTRIM PLUS 75% W/W WSP in drinking

water for 3 to 6 days. mg of product/kg live weight/day* x

average live

weight (kg) of poultry to
be treated

..... = mg of product per liter of
Daily water consumption (liter) Water
per animal

Half fill the container with potable water, add the prescribed amount of COSUTRIM PLUS, shake vigorously, then add the rest of the water.

The daily amount should be added to drinking water so that the required medication is administered within 24 hours. Medicated drinking water and stock solutions should be replaced every 24 hours. Animals should not have access to other water sources during the period when medicated water is being administered to them. The dilution must be calculated so that the animals are sufficiently watered. At the end of treatment, care must be taken to clean the water bowl properly to prevent any ingestion of the active substance in sub-therapeutic quantities. Body weight and water intake should be determined as accurately as possible to ensure the correct dose is administered. Consumption of medicated water depends on the clinical condition of the animals. To obtain the correct dosage, the concentration of the veterinary medicinal product must be adjusted according

ROUTE OF ADMINISTRATION:

Oral

CONTRA-INDICATIONS

Do not use in animals hypersensitive or animals with liver or kidney disease.

Do not administer to animals with impaired hematopoietic function.

Do not use in case of known hypersensitivity to sulfonamides, trimethoprim, or any of the excipients.)

WARNINGS AND PRECAUTIONS**Special warnings for each target species**

Avoid under dosing. Sick animals may drink less water than normal. If animals can be isolated, the concentration of the veterinary medicine can be proportionally increased in the water.

Regular monitoring of water consumption is recommended, especially in broiler chickens

Special precautions for use

Special precautions for use in animals

Due to the variable sensitivity (in time and space) of bacteria to potentiated sulfonamides, the resistance of bacteria is likely to fluctuate from one region to another, or even from one farm to another. Therefore, it is recommended to take bacteriological samples and perform sensitivity tests. The administration of the veterinary medicinal product in question must be based on the culture and sensitivity of the microorganisms in the sick livestock animals or on the basis of recent previous experiences at the site concerned. If the veterinary medicinal product is used in ways other than those indicated in the summary of product characteristics, any risk of an increase in the prevalence of bacteria resistant to sulfachloropyridazine and trimethoprim and a reduction in the effectiveness of combinations of trimethoprim with other sulfonamides due to cross-resistance. During any use of the veterinary medicinal product, the official and local antimicrobial policy should be respected

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Sulphonamides may cause hypersensitivity (allergy) reactions after injection, inhalation, ingestion or contact with skin.

Hypersensitivity to sulfonamides may cause cross-reactions with other antibiotics. Allergic reactions caused by these substances can be serious in some cases.

Avoid contact with this veterinary medicinal product if you are sensitive to sulfonamides.

If after exposure to the veterinary medicinal product, symptoms such as a rash appear, seek medical advice immediately and show the physician the package leaflet.

Put on a pair of waterproof latex or rubber gloves and a pair of protective glasses before using this veterinary medicinal product.

In case of eye contact, rinse with plenty of clean water and, in case of irritation, call a doctor.

In case of accidental ingestion, consult a doctor.

Wash contaminated hands and skin immediately after handling the veterinary medicinal product

INTERACTIONS WITH OTHER MEDICAMENTS:

Do not combine with other veterinary drugs

PREGNANCY AND LACTATION:

The safety of the veterinary medicinal product has not been established in cases of egg laying.

SIDE EFFECTS:

Sulphonamides may affect renal function (crystalluria, hematuria, urinary obstruction) and hematopoiesis (thrombocytopenia, anemia).

The frequency of adverse reactions is defined as follows:

- very common (adverse effects in more than 1 animal in 10 animals treated)
- common (between 1 and 10 animals out of 100 animals treated)
- uncommon (between 1 and 10 animals out of 1,000 animals treated)
- rare (between 1 and 10 animals out of 10,000 animals treated)
- very rare (less than one animal in 10,000 animals treated, including isolated cases)

SYMPTOMS AND TREATMENTS OVERDOSE:

The target species tolerate doses at least 5 times higher than the recommended dose which are administered for a treatment period at least three times the maximum recommended period, without any adverse effects having been demonstrated.

WITHDRAWAL PERIOD:

Poultry (Chicken)

Meat and offal: 7 hours

Eggs : Do not use in hens laying eggs intended for human consumption

STORAGE CONDITIONS:

Store below 30°C

Protect from light and moisture

Jauhkan dari kanak-kanak

keep away from children

Shelf Life as packaged for Sale

3 years

Shelf Life after first opening of container

7days

Shelf life after reconstitution or dilution

The solution is discarded after 24 hours

PACKING:

100gm powder packed in aluminium pouch

MANUFACTURED BY:**NAM PHARMA SDN. BHD**

5, Lebuhr Perusahaan Kelebang 11, Kawasan Perindustrian Igb,
31200 Chemor, Negeri Perak

NAME & ADDRESS OF PRODUCT REGISTRATION HOLDER

FIJILIAM ENTERPRISE SDN. BHD.No. 100 & 101, Taman AST,
Jalan Sungai Ujong,
70200 SEREMBAN NEGERI SEMBILAN, MALAYSIA

DATE OF REVISION OF PACKAGE INSERT:

Feb 2024