



MELOX 1.5MG/ML SUSPENSION

Description and Composition

MELOX 1.5MG/ML SUSPENSION is a pale yellow or yellowish with green tinge suspension. Each ml contains 1.5mg Meloxicam as its active ingredient and 2mg Sodium Benzoate as its preservative.

Pharmacodynamics

Meloxicam is a non-steroidal anti-inflammatory drug (NSAID) of the oxicam class which acts by inhibition of prostaglandin synthesis, thereby exerting anti-inflammatory, analgesic, anti-exudative and antipyretic effects. It reduces leukocyte infiltration into the inflamed tissue. To a minor extent it also inhibits collagen-induced thrombocyte aggregation. *In vitro* and *in vivo* studies demonstrated that meloxicam inhibits cyclooxygenase-2 (COX-2) to a greater extent than cyclooxygenase-1 (COX-1).

Pharmacokinetics

Absorption

Meloxicam is completely absorbed following oral administration and maximal plasma concentrations are obtained after approximately 4.5 hours. When the product is used according to the recommended dosage regime, steady state concentrations of meloxicam in plasma are reached on the second day of treatment.

Distribution

There is a linear relationship between the dose administered and plasma concentration observed in the therapeutic dose range. Approximately 97% of meloxicam is bound to plasma proteins. The volume of distribution is 0.3L / kg.

Metabolism

Meloxicam is predominantly found in plasma and is also a major biliary excretion product whereas urine contains only traces of the parent compound. Meloxicam is metabolised to an alcohol, an acid derivative and to several polar metabolites. All major metabolites have been shown to be pharmacologically inactive.

Elimination

Meloxicam is eliminated with a half-life of 24 hours. Approximately 75% of the administered dose is eliminated via faeces and the remainder via urine.

Indication

Alleviation of inflammation and pain in both acute and chronic musculo-skeletal disorders in dogs.

Recommended Dose

For oral use. To be administered mixed with food or directly into the mouth. Initial treatment is a single dose of 0.2mg meloxicam / kg body weight (equivalent to 1.3ml product / 10kg bw) on the first day. Treatment is to be continued once daily by oral administration (at 24-hour intervals) at a maintenance dose of 0.1mg meloxicam / kg body weight (equivalent to 0.67ml product / 10kg bw).

For longer term treatment, once clinical response has been observed (after ≥ 4 days), the dose can be adjusted to the lowest effective individual dose reflecting that the degree of pain and inflammation associated with chronic musculo-skeletal disorders may vary over time.

A clinical response is normally seen within 3-4 days. Discontinue after 10 days at the latest if no clinical improvement is apparent.

Shake well before use.

Avoid introduction of contamination during use.

Particular care should be taken with regard to the accuracy of dosing.

Route of Administration

For oral administration.

Contraindication

Do not use in pregnant or lactating animals.

Do not use in dogs suffering from gastrointestinal disorders such as irritation and haemorrhage, impaired hepatic, cardiac or renal function and haemorrhagic disorders.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients. Do not use in dogs less than 6 weeks of age.

Warning and Precautions

Special warnings for each target species

None

Special precautions for use

Special precautions for use in animals

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of renal toxicity.

This product for dogs should not be used in cats as it is not suitable for use in this species.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to nonsteroidal anti-inflammatory drugs (NSAIDs) should avoid contact with the veterinary medicinal product.

In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

This product can cause eye irritation. In case of contact with the eyes, immediately rinse thoroughly with water.

Interaction with Other Medicaments

Other NSAIDs, diuretics, anticoagulants, aminoglycoside antibiotics and substances with high protein binding may compete for binding and thus lead to toxic effects. The product must not be administered in conjunction with other NSAIDs or glucocorticosteroids. Pre-treatment with anti-inflammatory substances may result in additional or increased adverse effects and accordingly a treatment-free period with such veterinary medicinal products should be observed for at least 24 hours before commencement of treatment. The treatment-free period, however, should take into account the pharmacological properties of the products used previously.

Pregnancy and Lactation

The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Do not use in pregnant or lactating animals.

Side Effects

Typical adverse reactions of NSAIDs such as loss of appetite, vomiting, diarrhoea, faecal occult blood, lethargy and renal failure have very rarely been reported from post-marketing safety experience. Very rare cases of haemorrhagic diarrhoea, haematemesis, gastrointestinal ulceration and elevated liver enzymes have been reported from post marketing safety experience.

These side effects occur generally within the first treatment week and are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

If adverse reactions occur, treatment should be discontinued and the advice of a veterinarian should be sought.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

Symptoms and Treatment of Overdose

In the case of overdosage, symptomatic treatment should be initiated.

Storage Condition

Store below 30°C. Protect from light.

Shelf life

3 years. Discard 6 months after first opening.

Packing

30ml, 60ml, 120ml

Manufacturer & Product Registration Holder

Range Pharma Sdn Bhd, No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia.

Tel: 603-61568708 Fax: 603-61568707

Email: support@vetscareline.com

Revised date: 9.4.2026