

Homecare® Zenmox Suspension

400mg/10ml

Name and Strength of Active Substance(s):

Each 10ml contains:
Albendazole 400mg

Preservative(s):

Methyl Paraben 1.8mg/ml
Propyl Paraben 0.2mg/ml

Product Description:

White colour suspension with orange flavour.

Pharmacodynamics:

Mechanism of Action:

Albendazole is a benzimidazole carbamate with antiprotozoal and anthelmintic effects against intestinal and tissue parasites. Albendazole exhibits larvicidal, ovicidal and vermucidal activity, and is thought to exert its anthelmintic effect by inhibiting tubulin polymerisation. This causes the disruption of the helminth metabolism, including energy depletion, which immobilises and kills the susceptible helminth.

Pharmacokinetics:

Absorption:

In man, albendazole is poorly absorbed (less than 5%) following oral administration.

Distribution:

Following oral administration of a single dose of 400mg albendazole, the pharmacologically active metabolite, albendazole sulphoxide, has been reported to achieve plasma concentrations of 0.698 microgram/ml when taken with breakfast.

Metabolism:

Albendazole rapidly undergoes extensive first-pass metabolism in the liver, and is generally not detected in plasma.

Elimination:

The plasma half-life of albendazole sulphoxide is 8½ h. Albendazole sulphoxide and its metabolites appear to be principally eliminated in bile, with only a small proportion appearing in the urine. Elimination from cysts has been shown to occur over several weeks following high and prolonged dosing.

Special Patient Populations:

Elderly:

No specific studies have investigated the effect of age on albendazole sulphoxide pharmacokinetics.

Renal Impairment:

The pharmacokinetics of albendazole in patients with impaired renal function has not been studied.

Hepatic Impairment:

The pharmacokinetics of albendazole in patients with impaired hepatic function has not been studied.

Indication:

Treatment of single or mixed infestations of intestinal parasites.

Recommended Dosage:

For infestation of roundworm, whipworm, hookworm and pinworm:

Adults and children above 2 years old: One full bottle (10ml) as a single dose.
Children 1 to 2 years old: Take half the adult dose (5ml).

For infestation of threadworm and tapeworm:

Adults and children above 2 years old:
One full bottle (10ml) as a single dose once a day for 3 consecutive days.

Not to be used in children aged under 1 year.

Shake well before use.

Route of Administration:

For oral administration only.

Contraindications:

1. Albendazole must not be used during pregnancy.
2. Albendazole is contraindicated in patients with a known history of hypersensitivity to albendazole or other constituents of the dose forms.

Warnings and Precautions:

1. If you develop severe headache, nausea and vomiting, fits and/or problems with your vision after taking this medicine, consult a healthcare professional immediately.
2. This medicine may cause liver problems. Please consult a healthcare professional if you experience yellowing of the skin or eyes.
3. If you have been diagnosed with liver or kidney problems, seek advice from a healthcare professional before taking this medicine.
4. Avoid contact with the skin, eyes, mouth or nose, as albendazole suspension may cause irritation.
5. Women of childbearing age should begin treatment during the first 10 days of starting

Item Code	: 204XX draft 10 July 2024
PM Code	: Ins.P ZMSs
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: mm
CURSOR (ATTACHMENT 2)	: mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 1 / CC

L
260mm
±
2mm

W
110mm ± 2mm

- their period, or immediately after a negative pregnancy test, in order to avoid taking this medicine during early pregnancy.
6. If you are breastfeeding, please seek advice from a healthcare professional before using this medicine.
7. Should not be administered during confirmed or suspected pregnancy.

Interactions with Other Medicaments:

Caution should be taken when albendazole is administered concurrently with these medications. Coadministration with carbamazepine may decrease the efficacy of albendazole. Whereas, when it is administered concomitantly with cimetidine, dexamethasone and praziquantel, the plasma levels of the active metabolite of albendazole may increase.

Pregnancy and Lactation:

Pregnancy:

Albendazole must not be used during pregnancy.

Lactation:

Adequate human and animal data on use during lactation are not available. Thus, albendazole should not be used during breastfeeding unless the potential benefits are considered to outweigh the potential risks associated with treatment. Patients who are breastfeeding must seek advice from a healthcare professional prior to taking this medicine.

Side Effects:

The following convention is used for classification of adverse reaction frequencies:

Very common: $\geq 1/10$

Common: $\geq 1/100$ and $< 1/10$

Uncommon: $\geq 1/1,000$ and $< 1/100$

Rare: $\geq 1/10,000$ and $< 1/1,000$

Very rare: $< 1/10,000$

Immune system disorders:

Rare: Hypersensitivity reactions including rash, pruritus and urticaria.

Nervous system disorders:

Uncommon: Headache and dizziness.

Gastrointestinal disorders:

Uncommon: Upper gastrointestinal symptoms (e.g. epigastric or abdominal pain, nausea, vomiting) and diarrhoea.

Hepatobiliary disorders:

Rare: Elevations of hepatic enzymes.

Skin and subcutaneous tissue disorders:

Very rare: Erythema multiforme, Stevens-Johnson syndrome.

Please contact a healthcare professional if you notice any of these symptoms.

Symptoms and Treatment of Overdose:

Symptoms: Gastrointestinal symptoms may occur in albendazole overdose such as nausea, vomiting, diarrhoea and abdominal pain.

Treatment: Gastric lavage may be performed in first 2 to 3 hours after ingestion and other supportive and symptomatic treatment may be carried out if necessary.

Effects on Ability to Drive and Use Machine:

Adverse effects on the ability to drive or operate machinery have not been observed.

Storage Condition(s):

Store at temperature below 30°C.

Protect from light and moisture.

Keep the cap tightly closed.

Keep out of reach of children.

Jauhkan daripada capaian kanak-kanak.

Shelf Life:

2 years.

Dosage Forms and Packaging Available:

Bottle of 10ml.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7 & 12, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 10 July 2024

Item Code	: 204XX draft 10 July 2024
PM Code	: Ins.P ZMSs
Measurement(mm) $\pm$ 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: mm
CURSOR (ATTACHMENT 2)	: mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 1 / CC

L
260mm
 $\pm$
2mm

W
110mm $\pm$ 2mm