

Package Insert of Qilu Pharmaceutical (Inner Mongolia)-20%w/w Chlortetracycline Premix		Date of Revision of Package Insert: May 28, 2024
FOR VETERINARY USE ONLY		CONTROLLED MEDICINE
Product Name: Qilu Pharmaceutical (Inner Mongolia)-20%w/w Chlortetracycline Premix		
Active Substance: Chlortetracycline (as Chlortetracycline Calcium Complex)		
Strength: 20%w/w		
Product Description: Brown granule, not lumpy or moldy, no unpleasant odor.		
Pharmacodynamics: Tetracyclines generally act as bacteriostatic antibiotics inhibiting protein synthesis by reversibly binding to 30S ribosomal subunits of susceptible organisms thereby preventing binding to those ribosomes of aminoacyl transfer-RNA. Tetracyclines are believed to reversibly bind to 50S ribosomes and additionally alter cytoplasmic membrane permeability in susceptible organisms. In high concentrations, tetracyclines can inhibit protein synthesis by mammalian cells.		
Pharmacokinetics: Tetracycline is readily absorbed after oral administration to fasting animals. Bio-availabilities are approximately 60 – 80%. The presence of food or dairy products can significantly reduce the amount of tetracycline absorbed, with reductions of 50% or more possible. Tetracyclines as a class are widely distributed in the body, including to the heart, kidney, lungs, muscle, pleural fluid, bronchial secretions, sputum, bile, saliva, urine, synovial fluid, ascitic fluid, and aqueous and vitreous humor. Only small quantities of tetracycline and oxytetracycline are distributed to the CSF and therapeutic levels may not be attainable. While all tetracyclines distribute to the prostate and eye, doxycycline or minocycline penetrate better into these and most other tissues. Tetracyclines cross the placenta, enter fetal circulation and are distributed into milk.		
Target species: Pigs, Chickens and Cattle		
Indication and Direction for Use:		
Swine		
For reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by group <i>E Streptococci</i> susceptible to chlortetracycline. 50-100 grams per ton of feed.		
For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. 10 milligrams per pound of body weight per day. Feed approximately 400 grams per ton of feed, varying with body weight and feed consumption to provide 10 milligrams per pound of body weight per day. Feed for not more than 14 days.		
Swine (breeding)		
For the control of leptospirosis (reducing the instances of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. 400 grams per ton of feed. Feed continuously for not more than 14 days.		
Cattle (beef, under 700 pounds)		
For the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. 350 milligrams per head per day in feed.		
Cattle (beef)		
For the control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline. 350 milligrams per head per day in feed.		
Cattle (growing, over 400 pounds)		
For the reduction of the incidence of liver abscesses. 70 milligrams per head per day in feed.		
Cattle (beef, over 700 pounds)		
For the control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline. 0.5 milligrams per pound of body weight per day in feed.		
Cattle (calves, beef, non-lactating dairy)		
For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to chlortetracycline. 10 milligrams per pound of body weight per day. Feed approximately 400 grams per ton of feed, varying with body weight and feed consumption to provide 10 milligrams per pound of body weight per day. Treat for not more than 5 days.		
Chickens (broilers/fryer)		
For the control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. 100-200 grams per ton of feed. Feed continuously for 7 to 14 days.		
For the reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. 500 grams per ton of feed. Feed for 5 days.		
For the control of chronic respiratory disease and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. 200-400 grams per ton of feed. Feed continuously for 7 to 14 days.		
Do not feed to chickens producing eggs for human consumption.		
Route of Administration: Oral		
Contraindications: Chlortetracycline is contraindicated in patients hypersensitive to it or other tetracyclines. Because tetracyclines can retard fetal skeletal development and discolor deciduous teeth, they should only be used in the last half of pregnancy when the benefits outweigh the fetal risks. Oxytetracycline, chlortetracycline and tetracycline are considered more likely to cause these abnormalities than either doxycycline or minocycline. In patients with renal insufficiency or hepatic impairment, chlortetracycline must be used cautiously. Lower than normal dosages are recommended with enhanced monitoring of renal and hepatic function. Avoid concurrent administration of other nephrotoxic or hepatotoxic drugs. Because it may cause clostridial enterotoxemia in guinea pigs, chlortetracycline should not be used this species.		
Warnings and Precautions:		
Do not feed to chickens producing eggs for human consumption. Do not use in calves to be processed for veal.		
- Keep out of reach of children		
- Jauhkan daripada kanak-kanak		
Interactions with Other Medicaments: Tetracyclines may interfere with bactericidal activity of the penicillins, cephalosporins, and aminoglycosides. When orally administered, tetracyclines can chelate divalent or trivalent cations that can decrease the absorption of the tetracycline or the other drug if it contains these cations; it is recommended that all oral tetracyclines be given at least 1 – 2 hours before or after the cation-containing products.		
Statement on usage during pregnancy and lactation: Tetracyclines are excreted in milk, but because much of the drug will be bound to calcium in milk, it is unlikely to be of significant risk to nursing animals.		
Adverse Effects/Undesirable Effects: Chlortetracycline given to young animals can cause discoloration of bones and teeth to a yellow, brown, or gray color. High dosages or chronic administration may delay bone growth and healing.		
Overdose and Treatment		
Tetracyclines are generally well tolerated after acute overdoses. High oral doses given to ruminants, can cause ruminal microflora depression and ruminoreticular stasis.		
Withdrawal Period(s):		
Chickens: Withdraw 24 hours before slaughter when fed at 500 grams per ton. No withdrawal period when fed at 100-400 grams per ton when used according to label.		
Swine: Withdraw 5 days before slaughter at 10 milligrams per pound of body weight. No withdrawal period when fed at 50-400 grams per ton when used according to label.		
Cattle (beef, growing over 400 pounds): No withdrawal period when used according to label at 70 mg per head per day. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.		
Cattle (beef): Withdraw 48 hours prior to slaughter at 350 milligrams per head per day.		
Cattle (beef, under 700 pounds): Withdraw 48 hours prior to slaughter at 350 milligrams per head per day.		
Cattle (beef, over 700 pounds): Withdraw 48 hours prior to slaughter at 0.5 milligrams per pound of body weight per day.		
Cattle (calves, beef, non-lactating dairy): Withdraw 10 days prior to slaughter at 10 milligrams per pound of body weight per day. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.		
Shelf life: 2 Years		
Packaging: 25kg/bag		
Storage: Close tightly, avoid sunlight, keep in a dry place, store below 30°C (room temperature).		
Manufacturer Name: QILU PHARMACEUTICAL (INNER MONGOLIA) CO., LTD.		
Add: No.2 Wei Si Road, Jinchuannan District, Economy & Technology Development Zone, Hohhot City, China.		
Product register holder & distributed by: Shennong Animal Health (Malaysia) Sdn Bhd.		
Add: No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1, 47610 Subang Jaya, Selangor, Malaysia		