

PRODUCT LITERATURE

PRODUCT NAME

Amoxyle 50% w/w WSP

EACH GRAMS CONTAINS:

Active Ingredient:

Amoxicillin trihydrate 500mg

Product Description:

White Powder

Target Species:

Chickens, ducks, turkeys, pigs.

Pharmacodynamics:

Amoxicillin is a bacterial semisynthetic penicillin with a broad spectrum of activity against Gram positive and Gram negative bacteria. IT owes its activity to the inhibition of the development of the peptidoglycan network structure in the bacterial cell wall.

Pharmacokinetics:

Amoxicillin is well absorbed following oral administration and it is stable in the presence of gastric acids. Excretion of amoxicillin is mainly in the unchanged form via the kidneys to give high concentration in renal tissue and urine. Amoxicillin is well distributed in body fluids. Studies in birds have indicated that amoxicillin is distributed and eliminated more rapidly than in mammals. Biotransformation appeared a more important route of elimination in birds than in mammals.

Indication:

Treatment of infections in chickens, turkeys and ducks caused by bacteria susceptible to amoxicillin. Not effective against beta-lactamase producing organisms. Use of the product should be based on susceptibility testing and it should take into account official and local antimicrobial policies. Pigs: For the treatment of salmonellosis and pasteurellosis.

Recommended Dose:

Amounts to be administered and administration route Chickens, turkeys and ducks The product is administered in the drinking water. Prepare the solution with fresh tap water immediately before use. Once opened, use the contents of one sachet immediately. Any unused medicated water should be discarded after 12 hours. In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst being treated.

The following formula may be used to calculate the amount of product required per day (in grams):

$$\frac{\text{Number of birds} \times \text{average live weight (kg)}}{25 \text{ (for 20 mg/kg) or } 33 \text{ (for 15 mg/kg)}}$$

To ensure a correct dosage, bodyweight should be determined as accurately as possible to avoid underdosing. The uptake of medicated water depends on the clinical condition of the birds. In order to obtain the correct dosage the concentration of amoxicillin has to be adjusted taking into account water intake.

Chickens: The recommended dosage is 15 mg amoxicillin trihydrate per kg bodyweight. The total period of treatment should be for 3 days or in severe cases for 5 days.

Ducks: Recommended dosage is 20 mg amoxicillin trihydrate/kg bodyweight for 3 consecutive days.

Turkeys: Recommended dosage is 15-20 mg amoxicillin trihydrate/kg bodyweight for 3 days or in severe cases for 5 days.

Pigs: For the medication of Pigs, product may be administered in the drinking water or administered by addition to liquid feeds produced with commercial feed. It may not be used in dry feeds.

1. Administration via the drinking water

a. Administer in the drinking water to give 20mg/kg bodyweight daily. The dose should be divided and administered at approximately 12 hourly intervals for up to 5 days. Dissolve the contents of 150g of the product in the requisite quantity of water immediately before use.

b. For administration via the drinking water, the solution should be prepared with fresh potable water.

i. Any medicated water which is not consumed within 12 hours should be discarded.

ii. In order to ensure consumption of the medicated water, animals should not have access to other water supplies whilst on treatment.

1. Administration in liquid feed

a. Administer in the liquid feed, to give 20mg amoxicillin trihydrate/kg bodyweight daily for up to 5 days. Medicated feed should be freshly prepared on at least 3 occasions per day over the treatment period. The daily dose should be calculated based on the number of animals and average weight and then divided by the number of feed lots prepared in the day.

b. Medicated liquid feed should be prepared with fresh potable water.

c. Dissolve the required amount of product in water at a rate of 25g/L before addition to the feed. This may be done by dissolving the contents of 150g of the product in approximately 6L of water immediately before use. Alternatively, 67g of product which should be dissolved in approximately 2.5L of water immediately before use.

d. After adding the product to some or all of the water needed to make the liquid feed, ensure the product is fully dissolved. Dissolution of the product can take up to 10 minutes. This medicated water can then be mixed with the dry complete meal and if appropriate, the remaining water. The system used should ensure that the medicated water is evenly distributed into the feed. Once prepared the final medicated liquid feed should be fed to the pigs immediately.

e. The medicated liquid feed should not be fermented and should not be stored.

f. Stability of amoxicillin in all commercial feeds has not been established. In order to ensure that any loss of amoxicillin activity is minimized, the quantity of medicated liquid feed prepared should not exceed the amount of feed which will be consumed within 4 hours.

- i. Any medicated liquid feed which is not consumed within 4 hours should be discarded
- ii. Although restricted access to other water supplies would help ensure medicated liquid feed is consumed, separate clean potable water should remain available at all times for welfare reasons.

Contraindications:

AMOXYLE 50% W/W WSP should not be administered to rabbits, hamsters, gerbils and guinea pigs, or to birds producing eggs intended for human consumption.

Warning & Precautions:

Special warnings for each target species

None

Special precautions for use

- i. Special precautions for use in animals
Pigs: The uptake of medication by animals can be altered as a consequence of illness. In case of insufficient uptake of feed or water, animals should be treated parenterally.

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.
- ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa.
Allergic reactions to these substances may occasionally be serious.
 - 1) Do not handle this product if you know you are sensitised or if you have been advised not to work with such preparations.
 - 2) Handle this product with great care to avoid exposure, taking all recommended precautions.
 - 3) If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.
 - Avoid inhalation of dust. Wear either a disposable half-mask respirator conforming to European Standard EN149 or a non disposable respirator to European Standard EN140 with a filter to EN143
 - Wear gloves during preparation and administration of medicated water or liquid feed
 - Wash any exposed skin after handling the product or medicated water or feed

- Wash hands after use

Interactions With Other Medicaments:

None Known

Pregnancy & Lactation:

Laboratory studies in rats have not produced any evidence of a teratogenic effect due to the administration of amoxicillin.

Use only according to the benefit/risk assessment of the responsible veterinarian.

Side Effects:

None known

Symptoms and Treatment of Overdose

No problems with overdosage have been reported. Treatment should be symptomatic and no specific antidote is available.

Withdrawal Period

Chickens (meat & offal): 1 day

Ducks (meat & offal): 9 days

Turkeys (meat & offal): 5 days

Pigs (meat & offal): 2 days

Not for use in laying birds producing eggs for human consumption

Storage Condition

Store below 30°C protected from light

Shelf Life as packaged for Sale

2 years from the date of manufacturing

Shelf Life after first opening of container

To be used immediately upon opening the sachet

Shelf life after reconstitution or dilution

Medicated water: To be consumed within 12 hours

Medicated liquid feed: To be consumed within 4 hours

Disposal of Container

Disposed of in accordance with local requirements

Packing Size: Aluminum Pouch of 100gm, 500gm and 1kg.

Product Registration Holder/Distributed By:

GNC AGRITECH SDN. BHD.

24 Jalan Industri CheroK Tokun3, Taman Tokun Jaya,
14000 Bukit Mertajam, Pulau Pinang, Malaysia.

Manufactured By:

No6 & No8, Jalam CheroK To'kun2,
Taman Industri CheroK To'kun,
14000 Bukit Mertajam, Pulau Pinang, Malaysia.

Revision New

Date: Aug 2024