

VANTELIN GEL 1% w/w

Diclofenac Sodium 1%, the active ingredient of VANTELIN GEL 1% w/w, is a nonsteroidal drug that has anti-inflammatory and analgesic activity.

Ingredient(s):

Each gram contains:
Diclofenac Sodium 10mg

Route of administration:

Topical.

Pharmacodynamics:

ATC code: M02AA15

Mechanism of action and pharmacodynamic effects:

Diclofenac is a non-steroidal anti-inflammatory drug (NSAID) with effective analgesic, anti-inflammatory and antipyretic properties. Diclofenac exerts its therapeutic effects primarily through inhibition of prostaglandin synthesis by cyclo-oxygenase 2 (COX-2). VANTELIN GEL 1% w/w is an anti-inflammatory and analgesic preparation designed for topical application. In inflammation and pain of traumatic or rheumatic origin, VANTELIN GEL 1% w/w relieves pain and decreases swelling.

Pharmacokinetics:

Absorption

The quantity of diclofenac absorbed through the skin is proportional to the size of the treated area, and depends on both the total dose applied and the degree of skin hydration. Absorption amounts to about 6% of the applied dose of diclofenac after topical application of 2.5g diclofenac 1% gel on 500 cm² skin, determined by total renal elimination, compared with diclofenac tablets. A 10-hour occlusion leads to a three-fold increase in the amount of diclofenac absorbed.

Distribution

Diclofenac concentrations have been measured from plasma, synovial tissue and synovial fluid after topical administration of a diclofenac gel to hand and knee joints. Maximum plasma concentrations are approximately 100 times lower than after oral administration of the same quantity of diclofenac. 99.7% of diclofenac is bound to serum proteins, mainly albumin (99.4%). Diclofenac accumulates in the skin which acts as reservoir from where there is a sustained release of drug into underlying tissues. From the skin and underlying tissue, diclofenac preferentially distributes and persists in deep inflamed tissues (such as the joint), rather than in the bloodstream. Diclofenac is found in tissues at concentrations up to 20 times higher than in plasma.

Metabolism

The biotransformation of diclofenac involves single and multiple hydroxylation steps followed by glucuronidation, and glucuronidation of the intact molecule.

Elimination

The total systemic clearance of diclofenac from plasma is 263±56 ml/min. The terminal plasma half-life is 1-2 hours. Four of the metabolites, including the two active ones, also have short plasma half-lives of 1-3 hours. One metabolite, 3'-hydroxy-4'-methoxy-diclofenac, has a longer half-life but is virtually inactive. Diclofenac and its metabolites are excreted mainly in the urine.

Special populations

Renal and hepatic impairment

No accumulation of diclofenac and its metabolites is to be expected in patients suffering from renal impairment. In patients with chronic hepatitis or non-decompensated cirrhosis, the kinetics and metabolism of diclofenac are the same as in patients without liver disease.

Indication(s):

- For the external treatment of pain, inflammation and swelling in:
- injuries of the tendons, ligaments, muscles and joints, e.g. sprains, bruises, strains and back pain after sport or accidents;
 - localized forms of soft tissue rheumatism such as tendinitis (tennis elbow), shoulder-hand syndrome, bursitis, periarthropathies;
 - and for the symptomatic treatment of osteoarthritis of small and medium-sized joints located close to the skin such as the finger joints or knee joints.

Recommended Dosage:

For cutaneous use only.
Patients should consult their doctor if the condition does not improve or worsens within 7 days of starting treatment.

For adults and adolescents aged 12 years and over.

This medicine should be applied over the affected area 3 or 4 times daily and rubbed gently into the skin. The amount needed depends on the size of the painful area: 2 g to 4 g VANTELIN GEL 1% w/w (a quantity ranging in size from a cherry to a walnut) is sufficient to treat an area of about 400-800 cm². After application, the hands should be washed, unless they are the site being treated. The duration of treatment depends on the indication and clinical response. The product should not be used for more than 14 days.

Elderly patients (over 65 years of age)

The usual adult dosage may be used.

Children or adolescents (under 12 years of age)

The use and safety of VANTELIN GEL 1% w/w in children under 12 years has not been tested systematically and is not recommended.

Contraindications:

- Known hypersensitivity to diclofenac or to any of the excipients contained in VANTELIN GEL 1% w/w (see list of excipients).
- Patients in whom asthma, angioedema, urticaria, or acute rhinitis are precipitated by acetylsalicylic acid or other non-steroidal anti-inflammatory drugs (NSAIDs).
- During the last trimester of pregnancy.

Warnings and Precautions:

Special warnings and precautions for use

The possibility of experiencing systemic adverse events (those associated with the use of systemic forms of diclofenac) should be considered if topical diclofenac is used at a higher dosage or for a longer period of time than recommended (see Recommended Dosage). Topical diclofenac should be applied only to intact, non-diseased skin, and not to skin wounds or open injuries. It should not be allowed to come into contact with the eyes or mucous membranes, and should not be ingested. Discontinue the treatment if a skin rash develops after applying the product. Topical diclofenac can be used with non-occlusive bandages but should not be used with an airtight occlusive dressing.

Information concerning excipients

VANTELIN GEL 1% w/w contains propylene glycol and l-menthol, which may cause mild, localised skin irritation in some people.

Effects on ability to drive and use machines

Cutaneous application of VANTELIN GEL 1% w/w has no influence on the ability to drive and use machines.

Item Code	: 204xxxx - draft 26 Mar 2026
PM Code	: Ins.P VGO C
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: L000 - L000 mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 3 / CC

L
260mm
±
2mm

W
110mm ± 2mm

Statement on usage during pregnancy and lactation:

Pregnancy

There are insufficient data on the use of diclofenac in pregnant women. Diclofenac should be used during the first two trimesters of pregnancy only if the expected benefit justifies the potential risk to the fetus. As with other NSAIDs, use of diclofenac during the third trimester of pregnancy is contraindicated owing to the possibility of uterine inertia, fetal renal impairment with subsequent oligohydramnios and/or premature closure of the ductus arteriosus.

Lactation

It is not known whether topical diclofenac is excreted in breast milk. Diclofenac should only be used during lactation if the expected benefit justifies the potential risk to the newborn. If there are compelling reasons for using it, it should not be applied to the breasts nor should it be used at a higher dosage or for a longer period of time than recommended.

Interactions with Other Medicaments:

Since systemic absorption of diclofenac from topical application is very low, such interactions are very unlikely.

Side Effect(s) / Adverse Reaction(s):

Adverse reactions are listed below, by system organ class and frequency. Frequencies are defined as: very common (≥ 1/10); common (≥ 1/100 to < 1/10); uncommon (≥ 1/1,000 to < 1/100); rare (≥ 1/10,000 to < 1/1,000); very rare (< 1/10,000). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

<u>Infections and infestations</u> Very rare: Rash pustular.
<u>Immune system disorders</u> Very rare: Hypersensitivity (including urticaria), angioedema.
<u>Respiratory, thoracic and mediastinal disorders</u> Very rare: Asthma.
<u>Skin and subcutaneous tissue disorders</u> Common: Dermatitis (including contact dermatitis), rash, erythema, eczema, pruritus. Rare: Dermatitis bullous. Very rare: Photosensitivity reaction.

Overdose and Treatment:

The low systemic absorption of topical diclofenac renders overdose very unlikely. However undesirable effects, similar to those observed following an overdose of diclofenac tablets, can be expected if topical diclofenac is ingested (1 tube of 100 g contains the equivalent of 1 g diclofenac sodium). In the event of accidental ingestion, resulting in significant systemic adverse effects, general therapeutic measures normally adopted to treat poisoning with non-steroidal anti-inflammatory medicines should be used. Further management should be as clinically indicated or as recommended by the national poisons centres where available.

List of Excipients:

Propylene glycol, *l*-menthol powder, triethanolamine, isopropyl alcohol, Carbopol 940 and purified water.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in an airtight container. Store at temperature below 30°C Protect from light and moisture.

Product Description and Packing(s):

A colourless transparent gel with a pleasant refreshing odour.
Aluminium tube of 20g and 45g, with HDPE cap.

Date of revision:

12/Feb/2026

Item Code	: 204xxxx - draft 26 Mar 2026
PM Code	: Ins.P VGO C
Measurement(mm) ± 2mm	: 260(L) x 110(W)
CURSOR (ATTACHMENT 1)	: L000 - L000 mm
Colour of Artwork	: K 100
Source of Artwork	: MAC 3 / CC

L
260mm
±
2mm

Manufacturer:

Y.S.P. Industries (M) Sdn. Bhd. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13, Kawasan Perindustrian
Bandar Baru Bangi, 43000 Kajang, Selangor, Malaysia.

Product Registration Holder:

APD Pharma Sdn. Bhd. (201901008355)
No. 2-1, Jalan PDS 4, Telok Kemang, Jalan Pantai, PD Sunggala Gateway,
71050 Port Dickson Negeri Sembilan, Malaysia.

Product Owner:

Kowa Company, Ltd.
4-14, Nihonbashi-honcho 3-chome, Chuo-ku, Tokyo, JAPAN

Kowa Company, Ltd.

xxxxxx printing code xxxxxx

W
110mm ± 2mm