

CANYSTIN 4.4 % W/W PREMIX POWDER

Controlled Medicines/Ubat Terkawal

Product Description

A light brown color powder.

Composition

Each 1 gm contains

Active ingredients:

Nystatin..... 44 mg

Pharmacodynamics

Nystatin has a mechanism of action similar to that of amphotericin B. Nystatin binds to sterols in the membrane of the fungal cell altering the permeability of the membrane allowing intracellular potassium and other cellular constituents to leak "out". Nystatin has activity against a variety of fungal organisms, but is clinically used against topical, oropharyngeal and gastrointestinal candida infections.

Pharmacokinetics

Nystatin is not measurably absorbed after oral administration and almost entirely excreted unchanged in the faeces. The drug is not used parenterally because it is reportedly extremely toxic to internal tissues.

Indication

As an aid in the control of crop mycosis and mycotic diarrhea (candida albicans).

For the treatment of crop mycosis and mycotic diarrhea (candida albicans).

Target Species

Turkeys

RECOMMENDED DOSE

Aid in control of crop mycosis and mycotic diarrhea (Candida albicans). 50 grams per ton of feed.

For growing turkeys.

Treatment of crop mycosis and mycotic diarrhea

(Candida albicans). 100 grams per ton of feed.

For growing turkeys. To be fed for 7 to 10 days.

Route of Administration

Oral route by mixing into animal feed ration.

Contraindications

Nystatin is contraindicated for known hypersensitivity to it.

Warnings and Precautions

No special warnings are necessary for animal administration.

Interactions with Other Medicinal Products

No significant interactions reported for oral nystatin.

Pregnancy and Lactation

Although the safety of the drug during pregnancy has not been firmly established, the lack of appreciable absorption or case reports associating the drug with teratogenic effects appear to make it safe to use. It is not known whether nystatin is excreted in maternal milk, but because the drug is not absorbed after oral administration, it is unlikely to be of concern.

Side Effects

Occasionally high doses of nystatin may cause GI upset (anorexia, vomiting, diarrhea). Rarely, hypersensitivity reactions have been reported in humans.

Symptoms and Treatment of Overdose

Because the drug is not absorbed after oral administration, acute toxicity after an oral overdose is extremely unlikely, but transient GI distress may result.

Storage Condition

Store in a cool dry place at 30°C and below, away from direct sunlight.

Shelf Life as packaged for Sale

3 Years

Shelf life after opening

7 Days

Withdrawal Period

Growing turkey : Meat withdrawal time : 0 days

Packing:

25kg

Country Registration number:

For animal use only :

Manufactured by:

Nam Pharma Sdn. Bhd.

5, Lebuhr Perusahan Klebang 11,

Taman Perindustrian Antarabangsa IGB 312000,

CHEMOR, PERAK, MALAYSIA

Product Registration Holder:

Agro-farmchem Sdn Bhd, No. 5, Jalan Ss13/3a,

Subang Jaya Industrial Park,

47500 SUBANG SELANGOR MALAYSIA

Date of Revision: 30th October 2025