



KOTRA PHARMA (M) SDN BHD (90082-v)

AXCEL CINNARIZINE TABLET

AXCEL[®]
CINNARIZINE
TABLET

COMPOSITION:

Each tablet contains
Cinnarizine 25mg

PRESENTATION:

Light pink, biconvex, round tablets, 9mm in diameter with a single score-line on one side and marked KOTRA logo on the other side.

INDICATIONS:

Used for the symptomatic treatment of nausea and vertigo due to Meniere's disease and other labyrinthine disturbances and for the prevention and treatment of motion sickness. Cinnarizine is reported to possess smooth muscle relaxant properties and to inhibit vasoconstriction and is thus used in the management of various vascular disorders.

PHARMACOLOGY:

Mechanism of Action:

Cinnarizine is a histamine antagonists. Effectively antagonise the contraction of smooth muscle; the dilatation and increased permeability of the vasoconstrictor actions of histamine and vasodilator actions. It is a CNS depressant. Effective against motion sickness, relieve the nausea and vertigo due to labyrinthine disturbance.

Pharmacokinetics:

Cinnarizine are well absorbed from gastro-intestinal tract. It acts within about 30 minutes and its effect last for 4 hours. Sedative effects are not marked. It is metabolized in the liver and excreted mainly as metabolites in the urine.

DOSAGE & ADMINISTRATION:

For oral administration.

Adult and children over 12 years of age :

1 tablet, three times daily.

Children 5 - 12 years :

may be given half the adult dose.

Motion sickness :

Adults and children over 12 years : 1 tablet 2 hours before travelling and half tablet every 8 hours during the journey.

Children 5-12 years : half tablet 2 hours before travelling and 1/4 tablet every 8 hours during the journey.

CONTRAINDICATION:

The safe use of Cinnarizine in pregnancy has not been established and it should not therefore be administered to women who are pregnant. The drug should be used with reasonable caution in hypotensive patients.

PRECAUTION:

The preparation may cause drowsiness. Avoid driving or operating machinery. Avoid taking alcoholic drinks. Use is not recommended in pregnancy.

SIDE EFFECTS:

Rarely, allergic skin reactions and fatigue have been reported. Occasionally, drowsiness may occur. An aggravation or appearance of extrapyramidal symptoms has been reported extremely rarely in elderly people during prolonged therapy. The treatment should be reduced or stopped in such cases.

OVERDOSAGE AND TREATMENT:

There is no specific antidote to Cinnarizine and in the event of overdosage, gastric lavage is recommended.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

PACK QUANTITIES:

Available in blister pack of 10 x 10's and 100 x 10's.

Further information can be obtained from doctor, pharmacist or the manufacturer.

MAL19930647AZ

Product Registration Holder & Manufactured By :



Kotra Pharma (M) Sdn. Bhd.

No. 1, 2 & 3, Jalan TTC 12, (90082-V)

Cheng Industrial Estate,

75250 Melaka, Malaysia.

Date of revision : 10/11/22