

SWECON[®] SUSPENSION 100MG/ML

Ingredient(s):

Each ml contains:

Aluminium Magnesium Carbonate Hydroxide hydrate (Hydrotalcite) 100mg

Pharmacodynamics:

Swecon inhibits pepsin activity. Pepsin and other component of gastric juice do not affect the antacid action. Swecon acts locally and has no systemic effect.

Pharmacokinetics:

Swecon is not absorbed and does not interfere with the acid base balance. It has no laxative and no constipating effect and have been well-tolerate. Swecon acid-binding rate is 100% and acts very rapidly and persistently in the therapeutically optimal pH range between 3 and 5.

Indication(s):

Peptic ulcer, acute and chronic gastritis, hyperacidity.

Dosage and Administration:

Usual adult dose is 10ml, three to four times a day taken between meals or at bedtime, depending on the age and severity of the case.

Route of administration: Oral**Side Effect(s) / Adverse Reaction(s):**

Occasionally: Diarrhea, soft stool, dry mouth and anorexia.

Precaution(s) / Warning(s):

Swecon should be given with caution to patients with renal disorder, cardiac disorder, diarrhea and hypermagnesemia. Absorption may be decreased, when oral tetracyclines are used concurrently with antacids. Patients should be advised not to take antacid within 1 to 2 hours after administering tetracycline.

Symptoms and Treatment for Overdosage, and Antidote(s):

There is no specific antidote for its overdosage. Therefore, management of the patient consist of symptomatic and supportive therapy.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Shelf-Life:

3 years from the date of manufacture.

Product description:

A white color suspension

Packing(s):

Plastic container of 60ml, 100ml and 120ml.

Plastic container of 1L and 4L (for export only).

SWECON® SUSPensi 100MG/ML

Bahan aktif:

Setiap ml mengandungi:

Aluminium Magnesium Carbonate Hydroxide hydrate (Hydrotalcite) 100mg

Indikasi:

Ulser peptik, gastritis akut dan kronik, hiperasiditi.

Dos:

Untuk dewasa biasanya 10ml, 3 atau 4 kali sehari di antara masa makan atau sebelum tidur, bergantung kepada ketenatan.

適胃康懸液 100MG/ML

成份:

每錠含:

Aluminium Magnesium Carbonate Hydroxide hydrate (Hydrotalcite) 100mg

適應症:

最新合成之新型而理想之制酸劑，對消化性潰瘍，胃炎、胃酸過多等症有特效。

用法、用量:

成人每次服用10毫升，每日3-4次，於餐間或就寢前服用。



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P17, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679