

FORKNOW[®] TABLET

Forknow (Flunarizine) is a class IV calcium antagonist with a pharmacological profile which suggests its therapeutic potential in a number of neurological and cerebrovascular disorders.

Ingredient(s):

Each tablet contains:

Flunarizine (as Flunarizine Hydrochloride) 10mg

Pharmacodynamic:

Flunarizine is a class IV calcium antagonist which reduces the transmembrane fluxes of calcium in situations where calcium is stimulated to enter the cell in excess, thus preventing the deleterious consequences of "calcium overload" within the cell. It does not interfere with normal cellular calcium homeostasis having little negative inotropic effect on heart muscle and has no action on the myogenic tone of the blood vessels.

Pharmacokinetics:

1. Flunarizine is a selective calcium-entry blocker with a similar chemical structure and pharmacology profile to related compound, Cinnarizine. However, in contrast to Cinnarizine, it has long plasma half-life and needs only be given once a day. The majority of therapeutic trials in the prophylaxis of migraine, *occlusive peripheral, vascular disease* and vertigo of central peripheral origin have been placebo-controlled and have shown Flunarizine produces significantly greater beneficial effects than placebo as evaluated by subjective and objective criteria.
2. Flunarizine has also demonstrated vestibular depressive, antihistaminic, and anticonvulsant effects in human or animal.
3. Peak concentration levels occur 2 to 4 hours after oral administration of Flunarizine in healthy subjects. 0.8% of Flunarizine is present as free drug in plasma, 90% being bound to plasma proteins and 9% in blood cells. In healthy subjects Flunarizine is removed from the body with a long elimination half-life of around 18 days.

Indication(s):

For prophylaxis of migraine, peripheral venous insufficiency and occlusive arterial disease.

Dosage and Administration:

The usual dose for adults: 10mg tablet once daily preferably at night on retiring.

Route of Administration:

For oral administration.

Contraindication(s):

1. Patients known to be hypersensitive to this drug.
2. There is, as yet, no evidence of embryotoxic effects of Flunarizine in experimental animals. Nevertheless, the anticipated therapeutic benefits should be weighed against potential hazards before administering the compound during pregnancy.

Precaution(s) / Warning(s):

Flunarizine should be used with caution in patients with liver dysfunction and blood coagulating disorders.

Interactions with Other Medicaments:

When used in conjunction with anti-hypertensive drugs, dosage of the latter may need adjustment.

Galactorrhoea has been reported in some female patients on oral contraceptives within the first two months of flunarizine treatment.

Excessive sedation can occur when alcohol, hypnotics or tranquillisers are taken simultaneously with flunarizine.

The pharmacokinetics of flunarizine were unaffected by topiramate. After repeated dosing in migraine patients, systemic exposure to flunarizine increased by 14%. When flunarizine was co-administered with topiramate 50 mg every 12 hours, repeated dosing resulted in a 16% increase in systemic exposure to flunarizine. The steady-state pharmacokinetics of topiramate were unaffected by flunarizine. Chronic administration of flunarizine did not affect the disposition of phenytoin, carbamazepine, valproate or phenobarbital. Plasma concentrations of flunarizine were generally lower in patients with epilepsy taking these anti-epileptic drugs (AEDs) compared to healthy subjects given similar doses. The plasma protein binding of carbamazepine, valproate, and phenytoin is not affected by co-administration with flunarizine.

Pregnancy and Lactation:

Use during pregnancy

There are no data from the use of flunarizine in pregnant women. As a precautionary measure, it is preferable to avoid the use of flunarizine during pregnancy.

Use during lactation

It is unknown whether flunarizine is excreted in human milk. A decision on whether to discontinue breast-feeding or to continue/discontinue therapy with flunarizine should be made taking into account the benefit of breast-feeding to the child and the benefit of therapy to the woman.

Side Effect(s) / Adverse Reaction(s):

Asthenia, weight gain, migraine, drowsiness, depression, gastric pain, nausea, dry mouth, insomnia and rashes.

Effects on Ability to Drive and Use Machine:

Since somnolence may occur, especially at the start of the treatment, caution should be exercised during activities such as driving or operating dangerous machinery.

Symptoms and Treatment for Overdosage, and Antidote(s):

There is no specific antidote for its overdose, therefore, management of the patient consists of symptomatic and supportive therapy.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Shelf-Life:

3 years from the date of manufacture.

Product Description & Packing(s):

A slightly yellow color round tablet, one side impressed with a score and the other side with a mark "S"

Plastic bottle of 150's, 500's and 1000's (for export only)

Blister packing of 10's x 10

FORKNOW® TABLET

Bahan Aktif:

Setiap tablet mengandung:

Flunarizine (as Flunarizine Hydrochloride) 10mg

Indikasi:

Untuk profilaksis migrain, ketidak-cukupan vena periferall dan penyakit arteri oklusif.

Dos:

10mg satu kali sehari sebelum tidur.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679