

PRODUCT LITERATURE

KAOPEC SUSPENSION

Each 30 ml contains

Kaolin Light BP	9 gm
Pectin USP	200 mg

Preservative:

Methyl Paraben BP	0.1%w/v
Propyl Paraben BP	0.01%w/v

Product Description

Sweet taste, pink colour fruity and viscous suspension

Pharmacodynamics Properties

Mechanism of action/Effect:

Adsorbent and protectant. Kaolin is a natural hydrated aluminum silicate that is believed to adsorb large numbers of bacteria and toxins and reduce water loss. Pectin is a polyuronic polymer for which the mechanism of action is unknown. Pectin consists of purified carbohydrate extracted from citrus fruit or apple pomace. Studies have shown no decrease in stool frequency or fecal weight and water content with this combination even though stools appeared more formed.

Pharmacokinetics Properties

Absorption:

Not absorbed (up to 90% of pectin is decomposed in gastrointestinal tract).

Indication

For the relief of diarrhoea.

Recommended Dose

Adult: 4 - 8 tablespoon, after each loose stool
Child (above 6 years): 2 - 4 tablespoon, 4 times a day

Route of Administration

Oral

Contraindications

Risk-benefit should be considered when the following medical problems exist

- Dehydration - although adsorbent antidiarrheals may increase the consistency of feces and decrease the frequency of evacuation, they do not reduce the amount of fluid loss, but only mask its extent; rehydration therapy is essential if signs or symptoms of dehydration, such as dryness of mouth, excessive thirst, wrinkled skin, decreased urination, and dizziness or lightheadedness are present; fluid loss may have serious consequences, such as circulatory collapse and renal failure, especially in young children.
- Diarrhea, parasite-associated, suspected - use of adsorbent antidiarrheals may make recognition of parasitic causes of diarrhea more difficult; if parasitic agents are suspected pathogens,

appropriate stool analyses should be performed prior to therapy with adsorbents.

- Dysentery, acute, characterized by bloody stools and elevated temperature - sole treatment with adsorbent antidiarrheals may be inadequate; antibiotic therapy may be required.

Pregnancy and Lactation

Pregnancy/Reproduction

Pregnancy - Problems in humans have not been documented. Kaolin and pectin combination is poorly absorbed after oral administration.

Breast-feeding

Problems in humans have not been documented. Kaolin and pectin combination is poorly absorbed after oral administration.

Warnings and Precautions

Not recommended for children under 6 years of age

- Severe constipation, which may lead to faecal impaction, may rarely occur in children and the elderly patients taking kaolin and pectin. Kaolin and pectin may interfere with the absorption of other drugs, including antibiotics, administered concurrently.

PRECAUTION

Appropriate fluid and electrolyte therapy should be given to protect against dehydration. Oral rehydration therapy with the use of appropriate fluids including oral rehydration salts - remains the most effective treatment for dehydration due to diarrhoea. The intake of as much of these fluids as possible is therefore imperative.

Pediatrics

In infants and children up to 3 years of age with diarrhea, use is not recommended unless directed by a physician because of the risk of fluid and electrolyte loss. Oral rehydration therapy is recommended in children with diarrhea to prevent loss of fluids and electrolytes.

Geriatrics

In geriatric patients with diarrhea, caution is recommended because of the risk of fluid and electrolyte loss; these patients should be referred to a physician.

Interactions with Other Medicaments

Combinations containing any of the following medications, depending on the amount present, may also interact with this medication.

- Anticholinergics or other medications with anticholinergic activity
- Antidyskinetics
- Digitalis glycosides
- Lincomycins
- Loxapine
- Phenothiazines
- Thioxanthenes

- Concurrent use with kaolin and pectin combination may impair absorption of these medications when it is administered orally, resulting in decreased therapeutic effectiveness; it is recommended that kaolin and pectin combination be administered not less than 2 hours before or 3 to 4 hours after oral lincomycins; patients receiving digitalis should be monitored closely for evidence of altered effect.

Oral medications, other (prolonged use of adsorbents may interfere with absorption of other oral agents administered concurrently; it is recommended that kaolin and pectin combination be administered at least 2 to 3 hours before or after other oral medications).

Side Effects

Those indicating need for medical attention only if they continue or are bothersome

Incidence dose-related

- **Constipation** - usually mild and transient, but may rarely lead to fecal impaction.

Symptoms and Treatment of overdose

None

Packing

Packed in plastic container of 100 ml & 120 ml

Pharmaceutical Precautions

Store below 30°C in a dry place, protected from direct light.

Keep away from reach of children.

SHAKE WELL BEFORE USE.

Expiry Date

2 years from date of manufacturer

Name and Address of Product Registration Holder/ Distributor:

ZONTRON PHARMACEUTICALS SDN.BHD.
(445695-T)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Name and Address of Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Malaysian Drug Registration No. MAL19940173XC

Date of Revision: 16/10/2019