

PRODUCT LITERATURE

U-ALKALIZER POTASSIUM CITRATE MIXTURE

Each 10 mL contains:

Potassium Citrate 3 g

Preservative:

Methyl Paraben 0.12% w/v

Propyl Paraben 0.012% w/v

Description

Colourless, clear to slightly cloudy syrup with lemony odour and salty taste.

Pharmacodynamics Properties

Citrate and citric acid solutions are systematic and urinary alkalizers thereby providing symptomatic relief of dysuria.

Pharmacokinetic Properties

Potassium Citrate is absorbed and the citrate is metabolised to bicarbonate. Citric acid is metabolised to carbon dioxide and water. Oxidation is virtually complete with less than 5% of citrate being excreted unchanged in the urine.

Indication

For systemic or urinary alkalisation

Route of administration

Oral.

Recommended Dose

To be taken 3 times daily:

ADULT: 10-20 mL

CHILDREN up to 1 year: 2.5 mL,

1 - 5 years: 5 mL,

6 - 12 years: 10 mL

Directive:

Should be taken orally after meals, each dose can be diluted with drinking water (about 50 mL) before consuming.

Contraindication

Contraindicated in hyperkalaemia, renal dysfunction, ventricular arrhythmics and Addison's Disease.

Warning and Precautions

Potassium Citrate Mixture provides symptomatic relief only and is not anti-bacterial.

Effective anti-bacterial therapy should be co-prescribed. It should be used with caution when renal or cardiac dysfunction is present.

Caution must be exercised in renal or adrenal insufficiency. High doses of potassium may affect cardiac function.

Interactions with Other Medicament

Concurrent administration of potassium-containing drugs, potassium sparing diuretics or other drugs that increase potassium levels (e.g. ACE inhibitors, ciclosporin, aliskiren) may lead to hyperkalaemia. May interact with cardiac glycosides. Citrates alkalise the urine and thus may alter the urinary excretion of a number of drugs. This may lead to increased renal clearance of acidic drugs, such as salicylates, tetracyclines and barbiturates, and prolongation of the half-life of basic drugs, such as sympathomimetics and stimulants. Particularly noteworthy is the diminished anti-bacterial activity of nitrofurantoin and methenamine.

Pregnancy and Lactation

No adverse effects are anticipated at recommended doses when used for initial symptomatic relief only. Treatment with potassium citrate mixture is adjunctive and secondary to antibacterial treatment of urinary tract infection.

Side Effects

Mild nausea and occasionally vomiting may occur due to gastric irritation. Other side effects are those due to hyperkalaemia (if this occurs).

Symptoms and Treatment of Overdose

Overdosage is accompanied by nausea, vomiting, abdominal pain and symptoms due to hyperkalaemia and metabolic acidosis. Fluid and electrolyte balance together with ECG should be closely monitored.

Treatment is symptomatic and supportive. Moderate to severe hyperkalaemia is a medical emergency requiring prompt correction.

Packing

Packed in plastic container of 120 mL

Storage Conditions

Store below 30°C in a dry place, protected from direct light. Keep away from reach of children.

SHAKE WELL BEFORE USE.

Expiry Date

2 years from date of manufacture

Name and Address of Product Registration Holder/

Manufacturer:

TERAPUTICS SDN.BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Name and Address of Distributor:

ZONTRON PHARMACEUTICALS SDN. BHD (445695-T)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Malaysian Drug Registration No. :

MAL19940179XC

Date of Revision:

10/11/2021