

PRODUCT LITERATURE

BERIBION B1 TABLET 10MG

Each tablet contains

Thiamine Hydrochloride 10 mg

Preservative

Sodium Benzoate BP 0.1%w/w

Product description

Round, white coloured tablet.

Pharmacodynamics properties

Pharmacotherapeutic Group: Vitamin B1.

Thiamine is an essential co-enzyme for carbohydrate metabolism.

Pharmacokinetic properties

Thiamine is well absorbed from the gastrointestinal tract following oral administration, although the absorption of large doses is limited. It is widely distributed to most body tissues and appears in breast milk. Within the cell, thiamine is mostly present as the diphosphate. Thiamine is not stored to any appreciable extent in the body; amounts in excess of the body's requirements are excreted in the urine as unchanged thiamine or metabolites.

Indication

Vitamin B1 deficiencies.

Recommended dose

As prescribed by physician

Route of administration

Oral

Contraindication

Hypersensitivity to Thiamine Hydrochloride

Warnings and Precautions

Paediatrics

Problems in pediatrics have not been documented with intake of normal daily recommended amounts.

Geriatrics

Problems in geriatrics have not been documented with intake of normal daily recommended amounts. Studies have shown that the elderly may have impaired thiamine status, thereby requiring thiamine supplementation.

Interactions with Other Medicaments

The thiamine antagonists thiosemicarbazone and 5 fluorouracil can neutralise the effect of thiamine. Patients using any of these treatments may need their thiamine dose adjusted.

Thiamine could give false positive results for urobilinogen determination by the Ehrlich's reaction. High doses of thiamine may interfere with spectrophotometric assays of theophylline plasma concentration.

Pregnancy and Lactation

This product is not intended for use in pregnant or lactating women.

Side effects

Gastrointestinal disorders:

Mild gastrointestinal events such as nausea, vomiting, diarrhoea, and abdominal pain have been reported. Frequency not known (cannot be estimated from data).

Immune system disorders:

Hypersensitivity reactions have been reported (mainly after parenteral administration). Allergic and anaphylactic reactions, with symptoms of pruritus, urticaria, itching, hives, angioedema, abdominal pain, respiratory distress, tachycardia, palpitations, and shock have been reported in single cases. Frequency not known (cannot be estimated from data).

Symptoms and Treatment of Overdose

None

Packing

Packed in plastic container of 30 and 100 tablets

Storage Condition

Store below 30°C, in a dry place, protected from direct light.

Keep away from reach of children

Shelf life

2 years from date of manufacture.

Name and Address of Product Registration Holder/ Distributor:

ZONTRON PHARMACEUTICALS SDN.BHD.

(445695-T)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Name and Address of Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,

Lorong IKS Simpang Ampat B,

14100 Simpang Ampat, S.P.S.,

Pulau Pinang, Malaysia

Malaysian Drug Registration No. MAL19940197XC

Date of Revision: 29/04/2020