

PRODUCT LITERATURE

QURVION C TABLET 100MG

Each tablet contains:

Ascorbic Acid BP 50 mg
Sodium Ascorbate BP 57 mg
(equivalent to Ascorbic Acid BP 50 mg)

QURVION C TABLET 250MG

Each tablet contains:

Ascorbic Acid BP 125 mg
Sodium Ascorbate BP 140 mg
(equivalent to Ascorbic Acid BP 125 mg)

Description

Qurvion C Tablet 100mg: Oval-shape, yellow coloured tablet.
Qurvion C Tablet 250mg: Round, orange coloured tablet.

Pharmacodynamics / Pharmacokinetics

N/A

Indication

Health supplement.

Route of administration

Oral.

Recommended Dose

Qurvion C Tablet 100mg – Adult: 2-3 tablets a day or as prescribed by physician after meals

Qurvion C Tablet 250mg – Adults: 1 or more tablet, up to 500 mg a day, or as prescribed by physician after meals

Contraindication

Hypersensitivity to the active substances or to any of the excipients.

Warning and Precautions

Pediatrics

Problems in pediatrics have not been documented with intake of normal daily recommended amounts.

Geriatrics

Problems in geriatrics have not been documented with intake of normal daily recommended amounts.

Dental

Excessive use of chewable ascorbic acid tablets has been reported to cause breakdown of tooth enamel.

Interactions with Other Medicament

No interaction studies have been performed.

Pregnancy/Reproduction

Vitamin C may be taken by pregnant and lactating women under the supervision of a doctor.

Side Effects

Those indicating need for medical attention

Incidence dose-related

Kidney stones, oxalate (side or lower back pain)

Note: Occasionally, prolonged doses of ascorbic acid in excess of 1 g per day have been reported to cause an increase in urinary oxalate, which may cause precipitation of oxalate stones in the urinary tract in patients with renal disease, especially those on hemodialysis, or in patients with a history of renal stones. However, studies have not found an increase in urinary oxalate formation with high doses of ascorbic acid.

Those indicating need for medical attention only if they continue or are bothersome

Incidence less frequent or rare

Dizziness or faintness - with rapid intravenous administration

With high doses

Diarrhea - with oral doses greater than 1 gram per day
flushing or redness of skin
headache
increase in urination, mild - with doses greater than 600 mg per day
nausea or vomiting
stomach cramps

Symptoms and Treatment of Overdose

No cases of overdose have been reported.

Packing

Packed in plastic container of 30's and 100's tablet.

Storage Conditions

Store below 30°C in a dry place, protected from direct light.
Keep away from reach of children.
Jauhi daripada capaian kanak-kanak.

Expiry Date

Qurvion C Tablet 100mg: 1 1/2 years from date of manufacture.
Qurvion C Tablet 250mg: 2 years from date of manufacture.

Name and Address of Product Registration Holder/ Distributor: ZONTRON PHARMACEUTICALS SDN.BHD. (445695-T)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Name and Address of Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Malaysian Drug Registration No. :

Qurvion C Tablet 100mg: MAL19940202XCZ
Qurvion C Tablet 250mg: MAL19940206XCZ

Date of Revision:

04/11/2019