

PRODUCT LITERATURE

EUSOL LOTION

Each 100 ml contains:

Solution A - Chlorine	0.25 g
Solution B - Sodium Acetate	136.00mg
Glacial Acetic Acid BP	0.6ml

Description

Solution A – Colourless, odour of chlorine and clear solution

Solution B - Colourless, odourless clear solution

Mixture of A + B - Colourless, odour of chlorine and clear solution

Pharmacodynamics / Pharmacokinetics

Data not Available

Indication

Used in cleaning wounds and aberrations on the body.

Direction/Application:

1. Empty all the content of bottle (B) into the bottle (A).
2. Close cap and mix well.
3. Use as indicated.

Route of administration

Topical

Recommended Dose

Wash the affected parts liberally with Eusol lotion 2 to 3 times a day

Contraindication

Data Not Available

Warning and Precautions

May dissolve blood clots and cause bleeding.

Chlorine gas is an irritant producing inflammation of the conjunctive and when inhaled causing pain and spasm of the larynx and bronchi. There is a feeling of burning and suffocation. Coughing occurs, and if high concentrations are inhaled syncope and almost immediate death may follow. The irritant effect of chlorine may produce extensive pulmonary oedema with cyanosis, venous engorgement, and rapid respiration. There may be vomiting and excitement; acidosis may develop. Death may follow within 24 hours from either circulatory failure of pulmonary oedema

Interactions with Other Medicament

Data not available

Pregnancy and Lactation

Data not available

Side Effects

May be irritating to skin

Inhalation of fumes causes coughing and choking and may cause severe respiratory tract irritation and pulmonary oedema.

Ingestion causes irritation and corrosion of mucous membranes with pain and vomiting oedema of the pharynx and larynx and (rarely) perforation of the esophagus and stomach. A fall in blood pressure, delirium and coma may occur

Symptoms and Treatment of Overdose

If spilled on skin, wash with copious amount of water.

If ingested, give water, milk or other demulcents. Antacids and sodium tiosulphate 1-2.5% solution may also be given.

Packing

Packed in plastic container of 100 ml

Storage Conditions

Store below 30°C in a dry place, protected from direct light.

Keep away from reach of children.

FOR EXTERNAL USE ONLY

Expiry Date

2 years from date of manufacture

Name and Address of Product Registration

Holder / Manufacturer:

TERAPUTICS SDN. BHD. (590500-W)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Name and Address of Distributor:

ZONTRON PHARMACEUTICALS SDN. BHD.
(445695-T)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B,
14100 Simpang Ampat, S.P.S.,
Pulau Pinang, Malaysia

Malaysian Drug Registration No. :

MAL19940222XC

Date of Revision:

31/03/2020