

## PRODUCT LITERATURE

### SCABICIDE CREAM

#### Each 100 g contains

Crotamiton 10% w/w

#### Preservative

Methyl Paraben 0.12% w/w

### SCABICIDE LOTION

#### Each 100 ml contains

Crotamiton 10% w/v

#### Description

Cream: White soft odourless cream.

Lotion: Clear colourless odourless lotion

#### Pharmacodynamic properties;

Pharmacotherapeutic group: other antipruritics and other ectoparasitocides, including scabicides. Crotamiton has a symptomatic action on pruritus and is an acaricide. As an acaricide agent, crotamiton is effective on the motor system of the mites by inducing irreversible cessation of spontaneous movements.

#### Pharmacokinetic properties:

Crotamiton penetrates rapidly into human skin. Low but measurable concentrations of crotamiton are found in plasma, with a maximum level after 4-10 hours, declining rapidly thereafter.

#### Indication

Treatment of scabies.

#### Recommended Dosage

Scabies:

##### Adults:

Rub into the skin of the whole body from the chin down, paying particular attention to all folds and creases. A second application is advisable 24 hours later. Clothing and bed linen should be changed the next morning.

A cleansing bath should be taken 48 hours after the last application.

##### Children:

In children under three years of age Scabicide should not be applied more than once a day.

Pruritus: Apply to the affected area 2-3 times daily until medication is completely absorbed. Scabicide will provide relief from irritation for 6-10 hours after each application.

**Scabicide can be used in children.**

#### DIRECTIONS OF USAGE:

After the patient has taken a warm bath, the skin should be well dried and Scabicide rubbed into the entire body surface (excluding the face and scalp) until no traces of the preparation remain visible on the surface.

The application should be repeated once daily, preferably in the evening, for a total of 3-5 days. Depending on the response, special attention should be paid to sites that are particularly susceptible to infestation by the mites (eg interdigital spaces, wrists, axillae and genitalia). Areas

where there is pus formation should be covered with a dressing impregnated with Scabicide. While the treatment is in progress the patient may take a bath shortly before the next application. After completion of the treatment, a cleansing bath should be taken followed by a change of bed linen and underclothing.

#### Route of Administration

Topical

#### Contraindications

Acute exudative dermatoses. Hypersensitivity to the active substance or to any of the excipients.

Risk-benefit should be considered when the following medical problems exist:

-Inflamed skin, acute

-Raw, oozing skin surfaces (condition may be exacerbated)

#### Warnings and Precautions

Children under three years of age usage should only be under medical supervision. Should not be used in buccal mucosa and in or around the eyes since contact with the eyelids may give rise to conjunctival inflammation. In case of accidental contact with the eyes, or buccal mucosa rinse thoroughly with running water. Should not be applied in the presence of exudative wounds, acute eczema, broken skin, or very inflamed skin. In the presence of eczematous scabies, eczema should be treated before the scabies. Scabicide cream should only be used in pregnancy, breast feeding or for genital itching under medical supervision.

Appropriate studies on the relationship of age to the effects of crotamiton have not been performed in the pediatric and geriatrics population. Safety and efficacy have not been established.

#### Interactions with Other Medicaments

No interaction studies have been performed.

#### Pregnancy and Lactation:

##### Pregnancy

There are no controlled studies in human pregnancy. Therefore is not recommended during pregnancy, especially in the first three months.

##### Breastfeeding

It is not known whether the active substance passes into breast milk after topical administration. Therefore mothers should not use whilst breastfeeding unless directed by a physician. If is used during breastfeeding it should not be applied to the nipple area.

##### Fertility

No data is available on the potential effects of crotamiton on fertility.

#### Side Effects / Adverse Reactions

The most commonly reported adverse reaction during treatment is pruritus. Contact dermatitis and hypersensitivity reactions like rash, eczema, erythema, skin irritation and angioedema may occur rarely.

#### Overdose and Treatment

In cases of accidental ingestion, acute intoxication symptoms may be observed such as nausea, vomiting and irritation of the buccal, oesophageal and gastric

mucosa. Rare cases of loss of consciousness and seizure were reported.

General measures to eliminate the drug and reduce its absorption should be undertaken.

Symptomatic treatment should be administered as appropriate. Moreover although very rare, risk of methaemoglobinaemia exists in case of accidental ingestion as well as in case of excessive cutaneous absorption. The symptoms usually disappear following the discontinuation of the drug, but in severe cases treatment with methylene blue may be considered.

#### **Storage**

Store below 30°C in a dry place, protected from direct light.

Keep away from reach of children.

**For external use only.**

#### **Shelf Life**

2 years from date of manufacture

#### **Packing**

Cream: Packed in plastic container cream jar of 50gm and 100 gm

Lotion: Packed in plastic bottle of 100 ml

#### **Name and Address of Product Registration Holder / Distributor:**

**ZONTRON PHARMACEUTICALS SDN. BHD.**  
(445695-T)

Lot 10 & 11, PERDA Industrial Park,  
Lorong IKS Simpang Ampat B,  
14100 Simpang Ampat, S.P.S.,  
Pulau Pinang, Malaysia.

#### **Name and Address of Manufacturer :**

**TERAPUTICS SDN. BHD.** (590500-W)

Lot 10 & 11, PERDA Industrial Park,  
Lorong IKS Simpang Ampat B,  
14100 Simpang Ampat, S.P.S.,  
Pulau Pinang, Malaysia.

#### **Malaysian Drug Registration No.:**

Cream: MAL19940583XCZ

Lotion: MAL19940584XCZ

Date of revision: 10/09/2024