



WORMEX SUSPENSION 250MG/5ML

Description: Yellow coloured, vanilla flavoured suspension.

Contents

Pyrantel pamoate USP equivalent to pyrantel base 250mg/5ml

Methylparaben 0.1%

Propylparaben 0.01%

Indications

Single and mixed infestation of pinworm, roundworm and hookworms.

Pharmacology

Pyrantel is employed as the pamoate salt. It is a depolarizing, neuromuscular blocking agent. It induces marked, persistent nicotinic activation which results in spastic paralysis of the worm. Pyrantel also inhibits cholinesterases in single muscle cells of the helminth. Pyrantel causes depolarization and increased spike discharge frequency, accompanied by tension. The helminths are then dislodged by peristaltic activity. Oral absorption is poor and incomplete and elimination by renal and faecal excretions

Adverse effects

Side effects occur only occasionally and are mild and transient, gastrointestinal disturbances, headache, dizziness, skin rashes and fever have been reported.

Precautions

Safety for use in pregnancy and lactation has not been established, Caution when used in patients with impaired liver function.

Drug interactions

Piperazine may antagonize the anthelmintic effects of pyrantel. Concurrent use is not recommended.

Dosage and Administration

Adult	: 2 teaspoonfuls as single dose. May be repeated in 2 - 3 weeks if required.
Children 2 - 6 years	: 1/2 - 1 teaspoonful as single dose.
7 - 12 years	: 1 - 2 teaspoonfuls as single dose.

Overdosage

Because of poor gastrointestinal absorption, pyrantel serum concentrations are low. However, if overdose occurs, early gastric lavage and supportive therapy are necessary. There is no specific antidote.

Storage	: Keep container tightly closed. Store in a cool, dry place below 30°C. Protect from light.
Presentation/ Packing	: 250mg / 5ml x 10ml
Shelflife	: 3 years
Manufactured & Distributed by	: Prime Pharmaceutical Sdn. Bhd. 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah, 14000 Bukit Mertajam, Penang, Malaysia.