Title: Wintinea Cream - PI Date: 28 Feb 2025

wintinea Cream
(Antifungal)

Active Ingredient:
Tolnaftate 1.0%w/w
Preservative - Chlorocresol 0.1%w/w

Presentations:
Tubes of 10g and 20g.

Product Descriptions:
White smooth cream.

Pharmacology:
Tolnaftate is a synthetic fungicidal agent for the treatment of superficial fungal infections of the skin. Its spectrum of action encompasses typical fungi responsible for various forms of tinea and pityriasis. Tolnaftate inhibits the growth of Epidermophyton, Trichophyton spp., Microsporum, Pityrosporum orbiculare by distorting the hyphae and stunts the mycelial growth of the fungi species.

Indications:
Wintinea Cream is indicated for the treatment of superficial fungal infection of the skin including athlete's foot (tinea pedis), jock itch (tinea cruris), body ringworm (tinea corporis) and pityriasis versicolor.

Directions of Use:
Wash and dry the infected area. Apply a small amount of the cream and rub into the infected area twice daily for 2 weeks. To prevent recurrence, continue treatment for another 2 weeks after symptoms disappear. Wash hands after use. A total of 4 to 6 weeks treatment period may be necessary. Longer treatment period may be required for nail infection to clear up. Treatment may be repeated if infection recurs.

Symptoms and Treatments of Overdose:
In case of accidental ingestion, seek medical advice or assistance immediately.

Contraindication:
Not suitable for people with known hypersensitivity to Tolnaftate.

Precautions:
For external use only. Avoid contact with eyes.
Discontinue use and consult physician if itching or burning do not improve within 10 days; become worse or irritation occurs.
Not recommended for nail or scalp infection.
Not recommended for use in infants and children up to 2 years of age except under the advice of a physician.
If symptoms persist, please consult a doctor.

Use in Pregnancy and Lactation:
It is not known whether topical application of Tolnaftate will be harmful to an unborn baby.
It is not known whether topical application of Tolnaftate passes through into breast milk.

Interactions with Other Medicaments:
No clinically significant drug interactions known. However, care should be taken when using together with other topical medicines as other topical medicines may affect the absorption or effectiveness of Tolnaftate application.

Undesirable Effects:
Wintinea Cream is essentially well tolerated when applied on the intact or broken skin, in either exposed or intertriginous areas. Rarely contact dermatitis, pruritus and other skin reactions may occur.

Storage:
Store below 30°C. Protect from light. Keep cap tightly closed.
Keep out of reach of children.

Shelf Life:
3 years from the date of manufacture.

Manufacturer and Product Registration Holder
WINWA MEDICAL SDN. BHD.
198101009677(75794-A)
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

MAL19961017X BRU20052848NP

Date of Revision: 28 February 2025 X.EWT-I.09

Krim wintinea
(Antifungus)

Ramuan Aktif:
Tolnaftate 1.0%w/w
Bahan Pengawet - Klorokresol 0.1%w/w

Saiz Bungkusan:
10g dan 20g tiub.

Deskripsi Produk:
Krim lembut berwarna putih.

Indikasi:
Krim Wintinea digunakan untuk merawat jangkitan fungus pada kulit termasuk kurap, panau, kurap air dan sengkela.

Cara Penggunaan:
Bersih dan keringkan bahagian kulit terjangkit. Sapukan selapis krim yang nipis pada kulit terjangkit 2 kali sehari selama 2 minggu. Untuk mengelakkan jangkitan berulang, teruskan rawatan selama 2 minggu selepas simptom reda. Cuci tangan selepas guna.
Rawatan 4 hingga 6 minggu mungkin diperlukan. Rawatan yang lebih lama diperlukan untuk penyembuhan jangkitan kuku. Rawatan boleh diulangi sekiranya jangkitan berlaku.

Perhatian:
Untuk kegunaan luaran sahaja. Elakkan dari terkena mata.
Sekiranya luka atau kegatalan tidak sembuh dalam masa 10 hari atau bertambah teruk, sila henti guna dan dapatkan nasihat doktor.
Tidak disyorkan pada jangkitan kuku atau kulit kepala.
Tidak disyorkan kepada bayi di bawah 2 tahun kecuali dinasihatkan oleh doktor.
Jika simptom berlarutan, sila dapatkan nasihat doktor.

Cara Penyimpanan:
Simpan di bawah 30°C. Jauhkan daripada cahaya. Tutupkan penutup dengan ketat.
Jauhkan daripada capaian kanak-kanak.

Tempoh Simpanan:
3 tahun dari tarikh pengilangan.

永良癣药膏

主要成份： 托萘酯 1.0%w/w
防腐剂 - 氯甲酚 0.1%w/w
包装 ：10 克及 20 克条装。
药剂样式： 白色软膏。
主治 ：永良癣药膏主要用于治疗香港脚、汗斑、股癣、表皮癣、花斑癣及皮肤之各种霉菌感染引起的病症。
用法 ：将患处洗净后抹干。然后将适量的药膏敷于患处，每日 2 次，持续 2 个星期。以防重患，痊愈后请再持续 2 个星期。用药后请洗手。
整个疗程需要大约 4 至 6 个星期。指甲感染需要更长的时间痊愈。
若感染重返，可持续治疗。
注意 ：只限于外用。避免触及眼睛。
若烧伤或瘙痒症在十天内没获得改善或持续恶化，请停止使用并延医诊治。
不宜使用于指甲或头皮感染的病症。
2 岁以下的孩童不宜使用，除非获得医生的指示。
如症状持续，请延医诊治。
贮存 ：置于摄氏 30 度以下，避光。确保把盖关紧。
储存在远离孩童的地方。
有效期限： 3 年(由制造日计起)。

Size: 75mm x 160mm
Dark Blue (Pantone 2767U)